

# Advanced Care Paramedic

## Medical Directives

ALS PCS 5.4



**Hamilton  
Health  
Sciences**

CENTRE FOR PARAMEDIC  
EDUCATION AND RESEARCH

2025 - V1

PRINT DATE 2025-06-02

Introduction

Airway/  
Breathing

Cardiac/  
Circulation

Level of  
Consciousness

Pain/Sedation/  
Nausea

Procedural

Palliative Care  
/ Research

Chemical  
Exposure


Medical  
References

Medication  
Information

Contact

Destination  
Guidelines

The Emergency Health Services Branch of the Ministry of Health Version 5.4 of the ALS Patient Care Standards will now be the standard of care. These standards and guidelines include significant advances to the paramedic scope of practice since they were last published. As the ALS PCS is a living document, this Medical Directive book may not be an accurate reflection of the current scope of practice and/or ALS PCS. Paramedics are to refer to the CPER website for access to the most up to date version of the ALS PCS and to their certification letter for currently authorized medications and procedures.



Dr. Paul Miller  
Regional Medical Director



Dr. Rupinder Singh Sahsi  
Assistant Medical Director



Dr. Ian Buchanan  
Assistant Medical Director

Note: This Paramedic guide contains content from the Ministry of Health Advanced Life Support Patient Care Standards, version 5.4 dated June 2, 2025. To access the full document please refer to [www.CPER.ca](http://www.CPER.ca).

© 2025 by the

Centre for Paramedic Education and Research  
430 McNeilly Road, Unit 201  
Stoney Creek, Ontario L8E 5E3  
Phone 905 521-2100 x71223  
Fax 905 643-1104

This material has been prepared and developed by the Centre for Paramedic Education and Research.

# Table of Contents

## 4 Introduction

### Airway/Breathing Medical Directives

- 21 Orotracheal Intubation
- 30 Supraglottic Airway
- 32 Bronchoconstriction
- 36 Moderate to Severe Allergic Reaction
- 38 Croup
- 41 Tension Pneumothorax
- 44 Continuous Positive Airway Pressure (CPAP)
- 46 Advanced Airway Suctioning & Reinsertion
- 49 Cricothyrotomy

### Cardiac/Circulation Medical Directives

- 53 Medical Cardiac Arrest
- 62 Trauma Cardiac Arrest
- 67 Newborn Resuscitation
- 73 Return of Spontaneous Circulation (ROSC)
- 77 Cardiac Ischemia
- 80 Acute Cardiogenic Pulmonary Edema
- 82 Cardiogenic Shock
- 84 Symptomatic Bradycardia
- 88 Tachydysrhythmia
- 95 Traumatic Hemorrhage

- 97 Hyperkalemia
- 100 Intravenous and Fluid Therapy
- 102 Adult Intraosseous
- 103 Pediatric Intraosseous

### Level of Consciousness Medical Directives

- 111 Hypoglycemia
- 116 Seizure
- 120 Opioid Toxicity and Withdrawal
- 123 Suspected Adrenal Crisis

### Pain/Sedation/Nausea Medical Directives

- 127 Analgesia
- 136 Procedural Sedation
- 138 Combative Patient
- 141 Nausea/Vomiting

### Procedural Medical Directives

- 145 Home Dialysis  
Emergency Disconnect
- 148 Emergency Childbirth
- 152 Central Venous Access  
Device Access
- 154 Lateral Patellar Dislocation

### Palliative Care / Research

- 157 Palliative Care Medical Directives
- 175 The PRIME Trial Medical Directive

## Table of Contents continued

### Chemical Exposure

- 181 Cyanide Exposure
- 183 Hydrofluoric (HF) Acid Exposure
- 185 Adult Nerve Agent Exposure
- 190 Pediatric Nerve Agent Exposure
- 195 Symptomatic Riot Agent Exposure

### Medical References

- 199 ETCO<sub>2</sub> Waveforms
- 200 12 Lead ECG Placement
- 201 STEMI Anatomical Location / 15 Lead ECG Placement
- 202 CPR Guidelines
- 204 Rule of Nines, Burn Percentage Chart
- 206 Intramuscular Injection
- 207 Formulas
- 208 Spinal Motion Restriction Standard Prompt Card
- 209 Dopamine Dosing Guide
- 210 Epinephrine 1 mg/mL = 1:1000 IM Dosing Guide
- 211 Morphine Dosing Guide
- 213 Fentanyl Dosing Guide
- 214 Seizure Medical Directive Dosing Guide

### 216 Medication Information

### 228 Contact

### Destination Guidelines

- 237 Field Trauma Triage Standards
- 240 Field Trauma Triage Prompt Card
- 241 Air Ambulance Utilization Standard
- 244 Deceased Patient Standards
- 245 Paramedic Prompt Card for Acute Stroke Protocol
- 246 Reporting to FACS Niagara
- 247 Paramedic Prompt Card for Sepsis
- 248 Paramedic Prompt Card for Sepsis (NEMS)
- 249 Niagara EMS Hospital Destination Policy
- 255 Canadian CSPINE Rule
- 256 STEMI Hospital Bypass Prompt Card
- 257 Patient Priority System (HPS)
- 259 GI Bleed Recognition Tool (HPS)
- 260 Isolated Hip Fracture Recognition Tool (HPS)
- 261 Psychiatric Emergency Recognition Tool (HPS)
- 262 Musculoskeletal Injury Recognition Tool (HPS)
- 263 Ebola Virus Disease Screening Recognition Tool (HPS)
- 264 Radio Channel Change Locations
- 265 FAST Sepsis Pre-Alert
- 266 STEMI Protocol Pearls
- 267 FIT2SIT - GWPS
- 269 List of Mandatory Patch Points



# Introduction

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



# Introduction

## ADVANCED LIFE SUPPORT PATIENT CARE STANDARDS

### Levels of Paramedics

In Ontario, there are 3 levels of qualification for paramedics which lead to Certification as a: Primary Care Paramedic (PCP), Advanced Care Paramedic (ACP), and Critical Care Paramedic (CCP). The qualification for each is set out in Ontario Regulation 257/00 made under the *Ambulance Act*, RSO 1990, c A-19. The qualifications for each include a requirement that the paramedic be authorized by a Medical Director of a Regional Base Hospital (RBH) to perform the controlled acts set out in Schedules 1, 2 and 3 to O. Reg. 257/00.

A paramedic may be authorized by the Medical Director to perform controlled acts from the Schedule immediately above their Certification. In this circumstance, the paramedic is required to perform the controlled act to a specific standard as set out in the *Advanced Life Support Patient Care Standards* (ALS PCS). All advanced medical procedures that are not listed as controlled acts in Schedules 1, 2 and 3, shall also be performed as set out in the ALS PCS.

### Purpose of Standards

The ALS PCS reflects current practices for paramedics in Ontario and provides benchmarks for paramedic performance. It also communicates the standards of practice and care by paramedics in Ontario to paramedics, patients, other disciplines and the public in general. In the provision of ALS PCS care, paramedics are required to ensure patient care and documentation is provided in accordance with all appropriate Standards as indicated in O. Reg. 257/00.

### Comprehensive Care

Although two patient care standards exist, both Standards represent a continuum of care that is to be followed in an integrated fashion during a call for service. While initiating and continuing treatment prescribed by these Medical Directives, a paramedic must ensure that the patient simultaneously receives care in accordance with the BLS PCS. It is acknowledged that there may be circumstances and situations where complying with ALS PCS is not clinically justified, possible, or prudent (e.g. multiple crews on scene, trapped patient, extenuating circumstances, competing patient care priorities). When treatment deviates from the standards, a paramedic must document the care provided, including reasoning for deviating from the ALS PCS.

## Format of the ALS PCS

This document is comprised of a Preamble section and six (6) sections: Section 1 – PCP Core Medical Directives; Section 2 – ACP Core Medical Directives; Section 3 – PCP Auxiliary Medical Directives; Section 4 – ACP Auxiliary Medical Directives; Section 5 – Certification Standard, and Section 6 – Research Trial Standard.

Airway /  
Breath.Cardiac /  
Circula.

## Use of the Medical Directives by Paramedics

These Medical Directives apply to paramedics who are authorized by a RBHP Medical Director to provide patient care. Delegation of controlled acts in the ALS PCS to paramedics falls under the exclusive oversight of the RBHP. Critical Care Paramedics and Advanced/Primary Care Flight Paramedics will perform controlled acts in accordance with the Base Hospital (RBHP) Medical Directives issued by the Ornge Base Hospital Medical Director(s).

LOC

Pain/  
Sed./  
Nausea

## Controlled Substances

**Please refer to the government of Canada's specific permissions granted for paramedics in Ontario under the *Controlled Drugs and Substances Act*.**

Proced.

### Counts

An inventory or 'count' shall be performed:

- When removing controlled substances from storage
- When returning controlled substances to storage
- When exchanging controlled between one paramedic to another

Pall Care /  
Research

A record of counts shall be maintained.

Chemical  
Exposure

### Storage/Transport

With the exception of performing counts, restocking or providing patient care, controlled substances shall be stored at all times either:

- On the person of a paramedic approved to administer or transport the controlled substances or,
- Secured by double locking (i.e. the controlled substances are contained in a locked pouch, bag, container, safe [or equivalent] which is locked inside a vehicle, room, mounted safe, mounted cupboard or equivalent)

Medical  
Refer.Medic.  
Info.

### Rationale

Ensures provincial compliance pursuant to subsection 56(1) of the Controlled Drugs and Substances Act (CDSA)

Contact

Destinat.  
Guide.

## General Structure of a Medical Directive

Airway /  
Breath.

All Medical Directives follow the same format and are comprised of the following sections:

Cardiac /  
Circula.

**Indications:** The general medical complaint or problem to which the Medical Directive applies.

LOC

**Conditions:** Clinical parameters that must be present for a procedure to be performed or for a medication to be administered.

Pain/  
Sed./  
Nausea

**Contraindications:** Clinical parameters that if present, preclude the performance of a procedure or the administration of a medication.

**Treatment:** Description of the type of procedure to be performed or the dosing of a medication.

Proced.

**Clinical Considerations:** Key clinical points that provide general guidance to the proper performance of a procedure or the administration of a medication.

Pall Care /  
Research

All of these sections must be taken into account before and during the implementation of a Medical Directive.

Chemical  
Exposure

## Auxiliary Medical Directives

Medical  
Refer.

Additional ("Auxiliary") controlled medical acts or advanced medical interventions may be delegated through use of the Auxiliary Medical Directives. Delegation of Auxiliary Medical Directives by a RBHP Medical Director to paramedics is optional and may be introduced after consultation and mutual agreement between the RBHP and the certified ambulance service that employs the paramedic. Some PCP and ACP Medical Directives contain the phrase, "(if available and authorized)". This phrase qualifies the skill or procedure as optional (*i.e.* auxiliary) even if included in PCP or ACP Medical Directives.

Medic.  
Info.

## Special Event Medical Directives

Contact

Medical Directives have been developed for time limited periods when a mass gathering could potentially strain the resources of the host community. These medical directives shall only be used by paramedics who have completed the necessary training and received Regional Base Hospital Program authorization.

Destinat.  
Guide.

## Consent to Treatment in Non-Emergency Situations

Except in emergency circumstances described below, paramedics shall obtain consent prior to administering treatment. If a patient is incapable of consenting to the treatment plan being proposed by a paramedic, consent may be given or refused on his or her behalf by the patient's substitute decision-maker (SDM). Consent may be expressed or implied. Implied consent may be assumed where a person provides a physical indication that they consent to the treatment plan being proposed. For example, a patient who cannot speak but extends his hand to a paramedic after the paramedic indicates she is going to perform a simple procedure, such as a blood glucose determination, may be giving implied consent to the treatment plan being proposed..

The elements required for consent to treatment are:

- consent must be given by a person who is capable of giving consent with respect to the treatment plan;
- consent must relate to the treatment plan;
- consent must be informed;
- consent must be given voluntarily; and
- consent must not be obtained through misrepresentation or fraud.

Consent to the treatment plan is informed if, before it is given by the person, he or she has:

- received the following information that a reasonable person in the same circumstances would require in order to make a decision about the treatment plan:
  - the nature of the treatment;
  - the expected benefits of the treatment;
  - the material risks of the treatment;
  - the material side effects of the treatment;
  - alternative courses of action;
  - the likely consequences of not having the treatment; and
- received responses to his or her requests for additional information about those matters.

Valid consent requires that a person has the capacity to provide consent. A person is presumed to have the capacity to provide consent with respect to

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Airway /  
Breath.

treatment and a paramedic may rely on that presumption unless the paramedic has reasonable grounds to believe that the person is incapable with respect to the treatment plan. A paramedic must perform a capacity assessment if it is not reasonable in the circumstances to presume the person is capable of consenting to the treatment plan.

Cardiac /  
Circula.

A patient is capable with respect to the treatment plan if the patient is:

LOC

- Able to **understand** the information that is relevant to making a decision about the treatment or alternatives being proposed; **and**
- Able to **appreciate** the reasonably foreseeable consequences of a decision or lack of decision with respect to the treatment plan.

Pain /  
Sed. /  
Nausea

If a patient is incapable of consenting to a proposed treatment plan, and the paramedic is aware or is made aware that the person has a prior capable wish with respect to the proposed treatment, they must respect that wish (for example, if the person does not wish to be resuscitated).

Proced.

Pall Care /  
Research

## Consent to Treatment in Emergency Situations

Where the person for whom the treatment is being proposed is apparently experiencing severe suffering or is at risk of sustaining serious bodily harm if the treatment is not administered promptly, it is considered to be an emergency.

Chemical  
Exposure

For situations involving consent to treatment in emergency situations, a paramedic shall comply with the applicable directions contained in the *Basic Life Support Patient Care Standards* (BLS PCS).

Medical  
Refer.

## Discharge from Care

Medic.  
Info.

If a paramedic is certified and authorized by their Regional Base Hospital to perform a prehospital discharge from care as per the applicable Medical Directives, the following applies. For the purpose of the applicable Medical Directives, a patient or substitute decision maker (SDM) present at the scene, must be capable to make an informed decision about their treatment plan.

Contact

Destinat.  
Guide.

**A paramedic authorized to perform a prehospital discharge from care shall:**

1. Determine whether a patient may be treated in accordance with the Treat and Discharge component of the applicable Medical Directive,
2. Communicate a clinically reasonable differential diagnosis to the patient or SDM,
3. Discuss the following elements of a discharge treatment plan:
  - a. The clinical situation related to the most likely diagnosis and/or differential diagnoses,
  - b. The symptoms and signs alerting them to seek further medical care (i.e. clues that the condition is worsening or that the diagnosis may not be correct),
  - c. Instructions regarding modifications(s) of activities of daily living following the health event,
  - d. Where possible, provide additional contacts for follow up care,
  - e. Instructions to call 911 back if their condition worsens or recurs, and
4. Ensure the patient has the necessary support to follow a discharge treatment plan. These supports may include:
  - a. access to food,
  - b. access to transportation,
  - c. access to alternate health care follow up,
  - d. a safe place to stay,
  - e. responsible adult at the scene available to monitor the patient, and
  - f. consideration of other apparent patient vulnerabilities.

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.**Refusal of Treatment**

If a patient refuses treatment, either in whole or in part, a paramedic shall comply with the applicable directions contained in the BLS PCS.

**Intravenous (IV) Access and Therapy by Primary Care Paramedics**

There are 2 types of authorization for PCPs IV cannulation and therapy.

“PCP Assist IV” is authorization for a PCP to cannulate a peripheral IV at the request and under the direct supervision of an ACP. The patient must require a peripheral IV in accordance with the indications listed in the Intravenous and Fluid Therapy Medical Directive - Auxiliary. The ACP will perform all IV therapy in accordance with the Intravenous and Fluid Administration Medical Directive once intravenous access is obtained. PCPs authorized in PCP Assist IV are not authorized to administer IV therapy.

Intro	
Airway / Breath.	This authorization level can no longer be obtained and only those who have previously received this authorization may continue to practice at this level.
Cardiac / Circula.	"PCP Autonomous IV" is authorization for a PCP to independently cannulate an IV according to the Intravenous and Fluid Therapy Medical Directive – Auxiliary. PCPs authorized in PCP Autonomous IV are authorized to administer IV therapy according to applicable Medical Directives.
LOC	Authorization for each type shall meet the requirements established by the OBHG MAC.
Home Medical Technology and Novel Medications	
Pain/ Sed./ Nausea	As community care advances, new home medical technologies and novel medications are being introduced for home use by patients and caregivers trained in the care required. They are generally used by patients with complex medical histories who may require emergent interventions which are not described in, or aligned with, the BLS PCS or ALS PCS.
Proced.	A "home medical technology" is an external or internal mechanical device prescribed by a member of a regulated health profession for the purpose of treating a medical condition.
Pall Care / Research	A "novel medication" is a self/caregiver-administered medication prescribed by a member of a regulated health profession that is required to treat patients with generally rare and unusually complex chronic medical conditions which are often end stage. The medication may be self/caregiver-administered by any route into any part of the body.
Chemical Exposure	
Medical Refer.	A paramedic may accept the claim that a patient or caregiver has knowledge and training about the technology or medication encountered. A paramedic may only assist a patient or caregiver within the authorized paramedic skill set.
Medic. Info.	For unusual circumstances requiring interventions in the out of hospital setting, the RBH may create local training modules, treatment guidelines or medical directives
Patching	
Contact	A paramedic shall patch to the Base Hospital when:
Destinat. Guide.	a) a medical directive contains a mandatory provincial patch point; <b>OR</b>



- b) for situations that fall outside of these Medical Directives where the paramedic believes the patient may benefit from online medical direction that falls within the prescribed paramedic scope of practice; **OR**
- c) for consultation when, in the paramedics opinion the patient presentation or situation warrants and medical advice is required.

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

In cases where a treatment option requires the prior authorization by the BHP AND the BHP cannot be reached despite reasonable attempts by the paramedic to establish contact, a paramedic may initiate the required treatment without the requisite online authorization if the patient requires a critical, potentially life-saving, intervention and, in the paramedic's opinion, the intervention would otherwise apply. All patch failures must be reported in a timely manner to the RBHP in accordance with local policy and procedures. Paramedics should document the attempts to patch to the BHP on the Ambulance Call Report (ACR).

If a BHP directs a paramedic to perform an assessment or intervention that exceeds the paramedic's scope of practice, the paramedic must advise the BHP of such and notify the physician that they cannot comply with the direction as it exceeds their scope of practice. In such cases, a paramedic should ask the BHP to provide alternative direction.

## Incident Reporting

Paramedics shall adhere to their ambulance service policies and the *Ontario Ambulance Documentation Standards* (incorporated by reference in Ontario Regulation 257/00) for incident reporting. Paramedics shall also adhere to additional RBHP policies regarding reporting of clinical care incidents to the RBHP.

## Responsibility for Care

Each paramedic is equally responsible for patient care within their scope of practice. If the care exceeds a paramedics scope of practice, responsibility for that continued care shifts to the higher certified paramedic.

If there is any disagreement between paramedics, the Base Hospital physician may be contacted. It is expected that when an intervention has been performed, the paramedic most appropriate for that intervention will remain responsible for the patient.

The risks to the patient during transport should be assessed and discussed prior to transferring care from a higher to lower level of paramedic (e.g.: ACP to PCP), paramedics must alert the highest-level paramedic of any change of patient status at any time in the call.

When transferring care from one level of paramedic to another, paramedics shall provide:

- a) current CTAS level;
- b) a history of the patient's current problem(s) and relevant past medical history;
- c) pertinent physical findings;
- d) a summary of management at scene/en route;
- e) the patient's response to treatment, including most recent vital signs; and
- f) the reason for transfer in cases of inter-facility transfers.

The transfer of responsibility of patient care is a critical juncture along the clinical care continuum. When transferring patient care to another health care provider (e.g. nurse, physician, etc.), a paramedic must comply with the BLS PCS regarding such transfers.

## Research

Clinical research is fundamental to the practice of medicine and the development of safer, more effective treatment options for patients. At times, research protocols require temporary changes to patient care standards. Changes to patient care standards will be approved and introduced by the MOH.

## Patient Care Model

Any patient care model subject to The Patient Care Model Standard (PCMS) requires approvals and training as per the PCMS. Paramedics shall assess and provide treatment to all patients in accordance with the ALS PCS and BLS PCS when patients do not completely meet the specific parameters of approved Patient Care Models.

## Conventions

"Conventions" refers to a consistent application of terms throughout the Medical Directives based on definitions below.

The word 'consider' is used repeatedly throughout the Medical Directives. Where this word appears, it indicates that a paramedic shall initiate the treatment when the indications are first identified unless there is strong clinical rationale to withhold or delay treatment or other extenuating circumstances. A paramedic must document his or her justification for withholding treatment on the ACR.

Airway /  
Breath.Cardiac /  
Circula.

## Medication Doses and Administration

Unless specified within the medical directive, the number of recommended medication doses may be administered regardless of any previous administrations. When more than one route of medication administration is listed, clinical circumstances for each case should determine the final route chosen.

LOC

When more than one route of medication administration is listed, the order of preference for route of administration is from left to right. Clinical circumstances for each case should determine the final route chosen.

Pain/  
Sed./  
Nausea

Pediatric medication doses can vary slightly according to the source of expert opinion. The pediatric medication doses in the ALS PCS are the preferred doses. However, medication doses as determined by an up-to-date version of a widely accepted RBHP approved pediatric emergency tape (e.g. Broselow Tape) are an acceptable alternative. Use of a pediatric emergency tape shall be documented on the ACR when it is used to determine a pediatric medication dose.

Proced.

Medication doses may be calculated based upon weight or other factors and result in a fraction that cannot be measured accurately. In these cases, the medication dose delivered will be rounded to the closest dose that can accurately be measured

Pall Care /  
Research

## Age and Vital Signs

Chemical  
Exposure

The general age cut off between adults and pediatrics is 18 years (under 18 yrs. is generally considered a pediatric patient). There is a wide range of "normal" for vital signs in adults and especially pediatrics. As much as possible, ages for pediatrics and cut off points for vital signs have been kept consistent throughout the Medical Directives. However, clinical research and expert opinion have resulted in a number of exceptions which in each case has been deliberately chosen and is clearly noted in each Medical Directive. Age will be written as a number of hours, days, or years throughout the medical directives. There is a deliberate gap in the definition of normotension and hypotension in adults.

Medical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

## ADULTS

**Normotension** SBP  $\geq$  100mmHg

**Hypotension** SBP < 90 mmHg

**Heart rate** Heart rate is always in beats per minute according to a cardiac monitor when it is applied. In situations where a cardiac monitor is not indicated then the heart rate is equal to the pulse rate.

**Bradycardia** HR < 50 BPM

**Tachycardia** HR  $\geq$  100 BPM

**Tachypnea** RR  $\geq$  28 breath/min

## PEDIATRICS

Age	Respiratory Rate	Heart Rate
0-3 months	30-60	90-180
3-6 months	30-60	80-160
6-12 months	25-45	80-140
1-3 yr	20-30	75-130
6 yr	16-24	70-110
10 yr	14-20	60-90

**Normotension** SBP  $\geq$  90 mmHg + (2 x age in years)

**Hypotension** SBP < 70 mmHg + (2 x age in years)

**Weight (kg)** (age x 2) + 10

## HYPOGLYCEMIA

Age	Blood glucose level
<2 yr	< 3.0 mmol/L
$\geq$ 2 yr	< 4.0 mmol/L

## Level of Awareness (LOA):

The word 'altered' refers to a GCS that is less than normal for the patient.

## Commonly Used Abbreviations

The following abbreviations, in alphabetical order, appear in the Advanced Life Support Patient Care Standards:

### A

ACP	Advanced Care Paramedic
ALS	Advanced Life Support
ALS PCS	Advanced Life Support Patient Care Standards
ASA	Acetylsalicylic acid
AED	automated external defibrillation

### B

BHP	Base Hospital Physician
BLS PCS	Basic Life Support Patient Care Standards
BPM	Beats per minute
BVM	Bag-valve-mask

### C

CCP	Critical Care Paramedic
COPD	Chronic obstructive pulmonary disease
COWS	Clinical Opiate Withdrawal Scale
cm	Centimeter
CPAP	Continuous positive airway pressure
CPR	Cardiopulmonary Resuscitation
CTAS	Canadian Triage and Acuity Scale
CVA	Cerebral vascular accident
CVAD	Central venous access device

### D

DKA	Diabetic ketoacidosis
DNR	Do Not Resuscitate
DSED	Dual Sequential External Defibrillation

### E

ECG	Electrocardiogram
ED	Emergency Department
ETCO <sub>2</sub>	End tidal carbon dioxide

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Intro	ETT	Endotracheal tube
Airway / Breath.	<b>F</b>	
	FiO <sub>2</sub>	Fraction of inspired oxygen
Cardiac/ Circula.	<b>G</b>	
	g	Gram
	GCS	Glasgow Coma Scale
	gtts	Drops
LOC	<b>H</b>	
	H <sub>2</sub> O	Water
	HR	Heart rate
Pain/ Sed./ Nausea	Hx	History
	HF	Hydrofluoric Acid
	<b>I</b>	
Proced.	IM	Intramuscular
	IN	Intranasal
	IO	Intraosseous
	IV	Intravenous
Pall Care / Research	<b>J</b>	
	j	Joule
Chemical Exposure	<b>K</b>	
	kg	Kilogram
	<b>L</b>	
Medical Refer.	LOA	Level of awareness
	LOC	Level of consciousness
Medic. Info.	<b>M</b>	
	Max.	Maximum
	mcg	Microgram
	MDI	Metered dose inhaler
	mg	Milligram
Contact	Min.	Minimum
	min	Minute
Destinat. Guide.		

mL/kg	Milliliter per kilogram
mmHg	Millimeters of mercury
MOH	Ministry of Health
ms	Milliseconds

## N

N/A	Not applicable
NaCl	Sodium chloride
nare	Nostril
NEB	Nebulized
NPA	Nasopharyngeal airway
NSAID	Non-steroidal anti-inflammatory drug

## O

OBHG-MAC	Ontario Base Hospital Group - Medical Advisory Committee
OPA	Oropharyngeal airway

## P

PCP	Primary Care Paramedic
PEA	Pulseless electrical activity
PPV	Positive Pressure Ventilation
PO	by mouth/oral
PRN	as needed

## Q

q	every
---	-------

## R

RBHP	Regional Base Hospital Program
ROSC	Return of spontaneous circulation
RR	Respiratory rate

## S

SAED	Semi-automatic external defibrillation
SC	Subcutaneous
SL	Sublingual
SBP	Systolic blood pressure

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Intro	STEMI	ST-segment elevation myocardial infarction
Airway / Breath.	<b>T</b>	
	TBI	Traumatic brain injury
	TCP	Transcutaneous pacing
Cardiac / Circula.	TOP	Topical
	TOR	Termination of Resuscitation
	<b>U</b>	
LOC	URTI	Upper respiratory tract infection
	<b>V</b>	
Pain / Sed./ Nausea	VCD	Vector change defibrillation
	VF	Ventricular Fibrillation
	VT	Ventricular Tachycardia
Proced.	VSA	Vital signs absent
	<b>W</b>	
Pall Care / Research	WNL	Within normal limits
Chemical Exposure	Reference and Educational Notes	
Medical Refer.	<p>The RBHPs have created a companion document of reference and educational notes intended to assist paramedics in implementing these Medical Directives. This will facilitate regular updating of these notes without having to issue frequent changes to the standards. It is expected that paramedics have mastered the relevant information as part of initial training and certification and have maintained their knowledge through continuing education and self assessment and reflective practice. The reference and educational notes do not define a standard of care and is not a nested document to this standard; however, they should be considered useful in ensuring that an appropriate standard of care is met.</p>	
Medic. Info.		
Contact		
Destinat. Guide.		



# Airway/Breathing

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



## Orotracheal Intubation Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Need for ventilatory assistance or airway control;

#### AND

Other airway management is ineffective

### CONDITIONS

#### **lidocaine spray**

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Orotracheal  
Intubation

#### **Orotracheal Intubation**

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### **lidocaine spray**

Allergy or sensitivity to  
lidocaine

Unresponsive patient

#### **Orotracheal Intubation**

Age <50 years **AND**  
current episode of asthma  
exacerbation **AND**  
not in or near cardiac arrest.

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider topical **lidocaine** spray (to the hypopharynx) for orotracheal intubation when GCS is  $\geq 4$

	Route
	TOP
Dose	10 mg/spray
Max. dose	5mg/kg
Dosing interval	N/A
Max. # of doses	20 sprays

Consider **oro**tracheal intubation

With or without intubation facilitation devices. The maximum number of intubation attempts is 2.

Confirm **oro**tracheal tube placement

Method	Method
Primary	Secondary
ETCO <sub>2</sub> (Waveform capnography)	ETCO <sub>2</sub> (Non-waveform device)
	Visualization
	Ausculation
	Chest rise
	Esophageal detection device

## CLINICAL CONSIDERATIONS

- ▶ An intubation attempt is defined as insertion of the laryngoscope blade into the mouth for the purposes of intubation.
- ▶ Confirmation of orotracheal intubation must use ETCO<sub>2</sub> (Waveform capnography). If waveform capnography is not available or not working then at least 3 secondary methods must be used. Additional secondary ETT placement confirmation devices may be authorized by the local medical director.
- ▶ ETT placement must be reconfirmed immediately after every patient movement.



NOTE: Refer to page 27 for **Pediatric ET Tube Sizing Chart**



NOTE: Refer to page 199 for **ETCO<sub>2</sub> Waveforms**

Intentionally Left Blank

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank

Intentionally Left Blank

## Pediatric Oral Endotracheal Tube (ETT) Sizing Chart

Pediatric Endotracheal Tube Size & Depth Estimation			
Size for children $\geq$ 1year of age:		$\left( \frac{\text{Age in years}}{4} \right) + 4$	
Size for children < 1year of age:	Gestational age	Weight	ETT Size
	< 28 weeks	< 1kg	2.5
	28-34 weeks	1-2 kg	3.0
	34-38 weeks	2-3 kg	3.5
	Term infant	> 3 kg	3.5
	1-6 months	3-5 kg	3.5-4.0
	6-11 months	5-10 kg	4.0
	1 year	10-12 kg	4.0-4.5
Depth (to the teeth) for children > 2 years of age:		$\left( \frac{\text{Age in years}}{2} \right) + 12$	



## Endotracheal Intubation

In general, ETI should be considered only when BLS airway maneuvers have not proven successful in maintaining oxygenation/ventilation. Several studies have indicated worse outcomes when patients are intubated under pre-hospital conditions. Paramedics should always consider the detrimental aspects of performing ETI and reference the Medical Directives to determine the appropriate patient population that may benefit.

- ▶ Attempt basic maneuvers as needed: positioning, suctioning, pharyngeal airway insertion, and BVM IPPV in addition to application of 100% O<sub>2</sub>.
- ▶ Initiate cardiac monitoring, and pulse oximetry (if available).
- ▶ Consider administration of Xylometazoline or Lidocaine (refer to ETI Medical Directive).
- ▶ Pre-oxygenate the patient for 30-60 seconds with 100% O<sub>2</sub> (and IPPV, if required).
- ▶ Choose the appropriate size ETT and check the cuff.
- ▶ Intubate the trachea, confirm tube placement, and secure the tube (see *Intubation Confirmation Procedure*).
- ▶ If intubation is unsuccessful, stop and re-oxygenate to avoid hypoxia. The paramedic may repeat attempt (maximum of 2 attempts) and/or initiate immediate transport.

### CONSIDERATIONS

- ▶ Should the patient require sedation post ETI, refer to the Patient Sedation Medical Directive.
- ▶ Alternative airway adjuncts should be readily available for use in the event of failed intubation.
- ▶ Endotracheal tube must be consistently evaluated for displacement, with special attention paid after every patient movement. The ETCO<sub>2</sub> waveform capnography is critically important for real time monitoring of ventilation status, as well as endotracheal tube placement.


## ENDOTRACHEAL TUBE CONFIRMATION

- Immediately after intubation, correct tube placement must be confirmed (refer to ETI medical Directive).


<i>Primary Method</i>	<i>Secondary Method</i>
ETCO <sub>2</sub> (Waveform capnography)	ETCO <sub>2</sub> (Non-waveform device)
	Auscultation
	Esophageal detection device
	Chest rise

- At least one primary and two secondary ETT placement confirmation must be used as per the ETI Medical Directive.
- An ETCO<sub>2</sub> device (quantitative or qualitative) must be used for ETT placement confirmation. It should also be used every time the patient is moved to confirm the ETT has not been dislodged.

### ET Tube Confirmation with ETCO<sub>2</sub>



### ET Tube Displacement with ETCO<sub>2</sub>



- The appearance of a 4 phase capnography waveform post ETI is indication of proper tube placement.
- Loss of the 4 phase capnography is an indication that the ET tube may have become displaced.
- Numeric value >10 mmHg or numeric value gradually rising is evident of correct placement.
- Numeric value of ETCO<sub>2</sub> may be low or zero if arrest has been prolonged. Generally ETCO<sub>2</sub> should be  $\geq 4$  mmHg and should continue to rise during CPR.
- When an ET tube is inserted into the esophagus of a patient who recently ingested carbonated beverages the initial ETCO<sub>2</sub> may be normal but will rapidly decrease to < 4mmHg after a few ventilations.

## Supraglottic Airway Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Need for ventilatory assistance or airway control

#### AND

Other airway management is ineffective

### CONDITIONS

#### Supraglottic Airway

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Absent gag reflex

### CONTRAINDICATIONS

#### Supraglottic Airway

Airway obstructed by a foreign object

Known esophageal disease (varices)

Trauma to the oropharynx

Caustic ingestion

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT

Consider **supraglottic airway insertion**

The maximum number of supraglottic airway insertion attempts is 2

Confirm **supraglottic airway placement**

Method	Method
<i>Primary</i>	<i>Secondary</i>
ETCO <sub>2</sub> (Waveform capnography)	ETCO <sub>2</sub> (Non-waveform device)
	Auscultation
	Chest rise

## CLINICAL CONSIDERATIONS

- ▶ An attempt at supraglottic airway insertion is defined as the insertion of the supraglottic airway into the mouth.
- ▶ Confirmation of supraglottic airway must use ETCO<sub>2</sub> (Waveform capnography). If waveform capnography is not available or is not working, then at least 2 secondary methods must be used.

## Bronchoconstriction Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Respiratory distress

#### AND

Suspected bronchoconstriction

### CONDITIONS

salbutamol	EPINEPHrine	dexamethasone
AGE: N/A	AGE: N/A	AGE: N/A
LOA: N/A	WEIGHT: N/A	WEIGHT: N/A
HR: N/A	LOA: N/A	LOA: N/A
RR: N/A	HR: N/A	HR: N/A
SBP: N/A	RR: BVM ventilation required	RR: N/A
Other: N/A	SBP: N/A	SBP: N/A
	Other: Hx of asthma	Other: Hx of asthma <b>OR</b> COPD <b>OR</b> 20 pack-year history of smoking

### CONTRAINDICATIONS

salbutamol	EPINEPHrine
Allergy or sensitivity to salbutamol	Allergy or sensitivity to EPINEPHrine
dexamethasone	
Allergy or sensitivity to steroids Currently on PO or parenteral steroids	

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **salbutamol**

	Weight <25 kg		Weight ≥25 kg	
	Route MDI*	Route NEB	Route MDI*	Route NEB
<i>Dose</i>	Up to 600 mcg (6 puffs)	2.5 mg	Up to 800 mcg (8 puffs)	5 mg
<i>Max. single dose</i>	600 mcg	2.5 mg	800 mcg	5 mg
<i>Dosing interval</i>	5-15 min. PRN	5-15 min. PRN	5-15 min. PRN	5-15 min. PRN
<i>Max. # of doses</i>	3	3	3	3

\* 1 puff=100mcg

Consider **EPINEPHrine**

Concentration 1 mg/mL = 1:1,000	
Route IM	
<i>Dose</i>	0.01 mg/kg*
<i>Max. single dose</i>	0.5 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

**\*The EPINEPHrine dose may be rounded to the nearest 0.05 mg.**

Consider **dexamethasone**

	Route
	PO/IM/IV
<i>Dose</i>	0.5 mg/kg
<i>Max. single dose</i>	8 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

## CLINICAL CONSIDERATIONS

- ▶ EPINEPHrine should be the 1<sup>st</sup> medication administered if the patient is apneic. Salbutamol MDI may be administered subsequently using a BVM MDI adapter.
- ▶ Nebulization is contraindicated in patients with a known or suspected fever or in the setting of a declared febrile respiratory illness outbreak by the local medical officer of health.
- ▶ When administering salbutamol MDI, the rate of administration should be 100 mcg approximately every 4 breaths.
- ▶ A spacer should be used when administering salbutamol MDI.

## EPINEPHrine 1 mg/mL = 1:1000 IM Dosing Chart

*Dose (0.01 mg/kg) is rounded to the nearest 0.05mg  
Use a 1 mL syringe*

AGE	WEIGHT	DOSE (mg)	VOLUME (mL)
3 months	5 kg	0.05 mg	0.05 mL
6 months	8 kg	0.08 mg	0.10 mL
9 months	10 kg	0.10 mg	0.10 mL
1 year	12 kg	0.12 mg	0.10 mL
2 years	14 kg	0.14 mg	0.15 mL
3 years	16 kg	0.16 mg	0.15 mL
4 years	18 kg	0.18 mg	0.20 mL
5 years	20 kg	0.20 mg	0.20 mL
6 years	22 kg	0.22 mg	0.20 mL
7 years	24 kg	0.24 mg	0.25 mL
8 years	26 kg	0.26 mg	0.25 mL
9 years	28 kg	0.28 mg	0.30 mL
10 years	30 kg	0.30 mg	0.30 mL
11 years	32 kg	0.32 mg	0.30 mL
12 years	34 kg	0.34 mg	0.35 mL
13 years	36 kg	0.36 mg	0.35 mL
14 years	38 kg	0.38 mg	0.40 mL
Adult	50 kg	0.50 mg	0.50 mL

**Note:** Dosage administered can be calculated by the weight based calculation in the Medical Directive and/or by using the above chart. Administered dosage in the chart may be rounded to the nearest volume increment that can be accurately measured.



## Moderate to Severe Allergic Reaction Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Exposure to a probable allergen

**AND**

Signs and/or symptoms of a moderate to severe allergic reaction  
(including anaphylaxis)

### CONDITIONS

#### EPINEPHrine

AGE: N/A

WEIGHT: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: For anaphylaxis only

#### diphenhydramINE

AGE: N/A

WEIGHT:  $\geq 25$  kg

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### EPINEPHrine

Allergy or sensitivity to EPINEPHrine

#### diphenhydramINE

Allergy or sensitivity to diphenhydramINE

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **EPINEPHrine**

	Route
	IM
	Concentration
	1 mg/mL = 1:1,000
<i>Dose</i>	0.01 mg/kg*
<i>Max. single dose</i>	0.5 mg
<i>Dosing interval</i>	Minimum 5 min
<i>Max. # of doses</i>	2

\* The EPINEPHrine dose may be rounded to the nearest 0.05 mg.

Consider **diphenhydramINE** (if available)

	Weight ≥25 kg to <50 kg	Weight ≥50 kg
	Route IV/IM	Route IV/IM
<i>Dose</i>	25 mg	50 mg
<i>Max. single dose</i>	25 mg	50 mg
<i>Dosing interval</i>	N/A	N/A
<i>Max. # of doses</i>	1	1

## CLINICAL CONSIDERATIONS

EPINEPHrine administration takes priority over IV access.



NOTE: Refer to page 35 for **EPINEPHrine 1mg/mL = 1:1000 IM Dosing Chart**.

## Croup Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Current history of URTI

#### AND

Barking cough or recent history of a barking cough

### CONDITIONS

#### EPINEPHrine

AGE: ≥ 6 months to < 8 years

LOA: N/A

HR: <200 bpm

RR: N/A

SBP: N/A

Other: Stridor at rest

#### dexamethasone

AGE: ≥ 6 months to < 8 years

LOA: Unaltered

HR: N/A

RR: N/A

SBP: N/A

Other: For mild, moderate and severe croup

### CONTRAINDICATIONS

#### EPINEPHrine

Allergy or sensitivity to EPINEPHrine

#### dexamethasone

Allergy or sensitivity to steroids

Steroids received within the last 48 hours

Unable to tolerate oral medications

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **EPINEPH**rine

	Weight <10 kg	Weight ≥10 kg
	Route	Route
	NEB	NEB
	Concentration	Concentration
	1 mg/mL = 1:1,000	1 mg/mL = 1:1,000
<i>Dose</i>	2.5 mg	5 mg
<i>Max. single dose</i>	2.5 mg	5 mg
<i>Dosing interval</i>	N/A	N/A
<i>Max. # of doses</i>	1	1

Consider **dexamethasone**

	Age ≥ 6 months to < 8 years
	Route
	PO
<i>Dose</i>	0.5mg/kg
<i>Max. single dose</i>	8 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

## CLINICAL CONSIDERATIONS

N/A

## Croup Assessment

- ▶ Croup is an upper respiratory infection that is generally the result of a viral infection.
- ▶ It tends to occur in children aged 6 months to 3 years, and is most prevalent at the age of 2 years.
- ▶ It is characterized by swelling and irritation of the respiratory tract, and is often associated with a “barking style” cough.
- ▶ The severity of the symptoms can be characterized using the guideline below.
- ▶ Generally speaking, patients with moderate to severe croup should be considered for therapy as per the Medical Directive.

### WESTLEY CROUP SCORE:

This allows the severity of symptoms to be classified. Maximum score possible is 17.

	Score					
	0	1	2	3	4	5
<b>Inspiratory Stridor</b>	-	Audible with stethoscope	Audible without stethoscope	-	-	-
<b>Retraction</b>	-	Mild	Moderate	Severe	-	-
<b>Air entry</b>	Normal	Decreased	Severely decreased	-	-	-
<b>Cyanosis</b>	None	-	-	-	With agitation	At rest
<b>Conscious level</b>	Normal	-	-	-	-	Altered

- ▶ Score of 2-3: Indicates mild croup.
- ▶ Score of 4-7: Indicates moderate croup.
- ▶ Score of >7: Indicates severe croup.

## Tension Pneumothorax Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Pre-arrest or VSA

**AND**

Absent or severely diminished breath sounds on the affected side(s)

### CONDITIONS

#### Needle Thoracostomy

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: Hypotension or VSA

Other: N/A

### CONTRAINDICATIONS

#### Needle Thoracostomy

N/A

### TREATMENT


Consider **Needle Thoracostomy**

### CLINICAL CONSIDERATIONS

Needle thoracostomy may be performed at the 4<sup>th</sup> intercostal space anterior axillary line (preferred location) **OR** the 2<sup>nd</sup> intercostal space in the midclavicular line.

## Chest Needle Thoracostomy

- ▶ Chest needle thoracostomy is performed in the setting of suspected tension pneumothorax. Refer to the Tension Pneumothorax Medical Directive for indications, conditions and contraindications.
- ▶ Prepare equipment including appropriate sized needle and syringe.
- ▶ Identify the appropriate landmark
  - 4th intercostal space, anterior axillary line (preferred)
  - 2nd intercostal space, mid clavicular line



- ▶ Cleanse the needle insertion site using aseptic technique.
- ▶ Insert the needle with attached syringe into the appropriate intercostal space on the affected side along the superior aspect of the rib with the bevel facing toward the rib.
- ▶ Aspirate for air as you advance the needle into the thoracic cavity.
- ▶ Once free air has been aspirated, advance the needle 2 mm further to ensure bevel and catheter are through the chest wall and into the pleural space.
- ▶ Advance the catheter over the needle to the hub, hold onto the catheter and remove the needle/syringe. A rush of air *may* be heard when you remove the needle/syringe.
- ▶ Place the needle into the biohazard container.
- ▶ Secure the catheter in place with tape.
- ▶ Place the flutter valve mechanism over the needle, being careful not to bend the catheter.
- ▶ Frequently reassess the patency of the needle thoracostomy site and clinical status of the patient

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

## CONSIDERATIONS

- ▶ The primary indicator of a successful needle decompression will be the improvement in patient condition in both hemodynamic status and partial or complete relief of respiratory symptoms.
- ▶ In the setting of a tension pneumothorax, it is preferred to not provide positive pressure ventilation (eg: with BVM) unless absolutely required and move quickly to chest needle decompression.



## Continuous Positive Airway Pressure (CPAP)

### Medical Directive – *AUXILIARY*

*An Advanced Care Paramedic may provide the treatment prescribed in this auxiliary Medical Directive if authorized.*

#### INDICATIONS

Severe respiratory distress

#### AND

Signs and/or symptoms of acute pulmonary edema or COPD

#### CONDITIONS

##### CPAP

AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: Tachypnea

SBP: Normotension

Other:  $\text{SpO}_2 < 90\%$  or accessory muscle use

#### CONTRAINDICATIONS

##### CPAP

Asthma exacerbation

Suspected pneumothorax

Unprotected or unstable airway

Major trauma or burns to the head or torso

Tracheostomy

Inability to sit upright

Unable to cooperate

## TREATMENT

### Consider **CPAP**

<i>Initial setting</i>	5 cm H <sub>2</sub> O	Or equivalent flow rate of device as per RBHP direction
<i>Titration increment</i>	2.5 cm H <sub>2</sub> O	Or equivalent flow rate of device as per RBHP direction
<i>Titration interval</i>	5 min.	
<i>Max. setting</i>	15 cm H <sub>2</sub> O	Or equivalent flow rate of device as per RBHP direction

### Consider increasing **FiO<sub>2</sub>** (if available)

<i>Initial FiO<sub>2</sub></i>	50-100%
<i>FiO<sub>2</sub> increment (if available on device)</i>	SpO <sub>2</sub> <92% despite treatment and/or 10cm H <sub>2</sub> O pressure or equivalent flow rate of device as per RBHP direction
<i>Max FiO<sub>2</sub></i>	100%

### Confirm **CPAP pressure by manometer** (if available)

## CLINICAL CONSIDERATIONS

N/A

## Advanced Airway and Tracheostomy Suctioning & Reinsertion Medical Directive

*An Advanced Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Patient with endotracheal tube, SGA (with gastric suction port) or tracheostomy tube

#### AND

Airway obstruction or increased secretions

### CONDITIONS

Suctioning (ETT / Tracheostomy)	Suctioning through SGA Gastric Port (if available)
AGE: N/A	AGE: N/A
LOA: N/A	LOA: N/A
HR: N/A	HR: N/A
RR: N/A	RR: N/A
SBP: N/A	SBP: N/A
Other: N/A	Other: Known or suspected gastric secretions or emesis following placement of SGA
	Persistent difficult ventilation despite other efforts to improve ventilation

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

**Emergency tracheostomy reinsertion**

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Patient with an existing tracheostomy where the inner and/or outer cannula(s) have been removed from the airway **AND**Respiratory distress **AND**Inability to adequately ventilate **AND** Paramedics are presented with a tracheostomy cannula for the identified patient.**CONTRAINDICATIONS****Suctioning  
(ETT/Tracheostomy)**

N/A

**Emergency tracheostomy  
reinsertion**

Inability to landmark or visualize

**Suctioning through SGA  
Gastric Port (if available)**

N/A

**TREATMENT****Consider Suctioning (ETT/Tracheostomy)**

	Age < 1 year	Age ≥ 1 year to < 12 years	Age ≥ 12 years
<i>Dose</i>	Suction at 60-100 mmHg	Suction at 100-120 mmHg	Suction at 100-150 mmHg
<i>Max. single dose</i>	10 seconds	10 seconds	10 seconds
<i>Dosing interval</i>	1 minute	1 minute	1 minute
<i>Max. # of doses</i>	N/A	N/A	N/A

Consider **Suctioning through SGA Gastric Port (if available)**

	Age < 1 year	Age ≥ 1 year to < 12 years	Age ≥ 12 years
<i>Dose</i>	Suction at 60-100 mmHg	Suction at 100-120 mmHg	Suction at 100-150 mmHg
<i>Max. single dose</i>	N/A	N/A	N/A
<i>Dosing interval</i>	N/A	N/A	N/A

Consider **emergency tracheostomy reinsertion**

The maximum number of attempts is 2

**CLINICAL CONSIDERATIONS****ETT/Tracheostomy Suctioning:**

Pre-oxygenate with 100% oxygen.

In an alert patient, whenever possible, have patient cough to clear airway prior to suctioning.

**Suctioning of SGA with gastric suction port:**

When gastric secretions are not evident, consider other causes of difficult ventilation (e.g., improper device size, incorrect depth, lack of posterior/inferior pressure, or airway obstruction) prior to attempting SGA suctioning.

Once fluid clears or if no fluid appears after 15 seconds, turn off suction.

**Emergency Tracheostomy Reinsertion:**

A reinsertion attempt is defined as the insertion of the cannula into the tracheostomy. A new replacement inner or outer cannula is preferred over cleaning and reusing an existing one.

Utilize a family member or caregiver who is available and knowledgeable to replace the tracheostomy cannula.

## Cricothyrotomy Medical Directive – *AUXILIARY*

*An Advanced Care Paramedic may provide the treatment prescribed in this auxiliary Medical Directive if authorized.*

### INDICATIONS

Need for advanced airway management;

#### AND

Intubation AND supraglottic airway insertion unsuccessful or contraindicated;

#### AND

Unable to ventilate

### CONDITIONS

#### Cricothyrotomy

AGE:  $\geq 12$  years

LOA: Altered

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### Cricothyrotomy

Suspected fractured larynx

Inability to landmark

## TREATMENT

Consider **cricothyrotomy**.

Confirm **cricothyrotomy tube placement**

Method <i>Primary</i>	Method <i>Secondary</i>
ETCO <sub>2</sub> (Waveform capnography)	ETCO <sub>2</sub> (Non-waveform device)
	Ausculation
	Chest rise

Airway /  
Breath.Cardiac/  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

## CLINICAL CONSIDERATIONS

Confirmation of cricothyrotomy must use ETCO<sub>2</sub> (Waveform capnography). If waveform capnography is not available or not working, then at least 2 secondary methods must be used. Additional secondary Cricothyrotomy tube placement confirmation devices may be authorized by the local medical director.

Cricothyrotomy tube placement must be reconfirmed immediately after every patient movement.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank



# Cardiac/Circulation

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



## Medical Cardiac Arrest Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Non-traumatic cardiac arrest.

### PRIMARY CLINICAL CONSIDERATIONS

In the following settings, consider very early transport after a minimum of one analysis (and defibrillation if indicated) once an egress plan is organized:

- 1) pregnancy presumed to be  $\geq 20$  weeks gestation (fundus at or above umbilicus, ensure manual displacement of uterus to left);
- 2) known reversible cause of the arrest unable to be addressed.

For patients in refractory VF or pulseless VT, consider:

- 1) Double sequential external defibrillation (DSED) if authorized, **OR**
- 2) Vector change defibrillation (VCD) if DSED is unavailable or not authorized, **AND**
- 3) Transport following three (3) doses of DSED or VCD and three (3) rounds of epinephrine if they remain in VF or pulseless VT (or after 3rd consecutive defibrillation if no IV/IO/CVAD/ETT access).

Refractory VF or pulseless VT is defined for the purpose of this directive, as persistent VF or pulseless VT after 3 consecutive shocks.

## CONDITIONS

### manual defibrillation

AGE: ≥ 24 hours  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: VF **OR** pulseless VT

### AED or SAED Defibrillation

AGE: ≥ 24 hours  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: Defibrillation indicated

### EPINEPHrine

AGE: ≥ 24 hours  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: Anaphylaxis suspected as causative event, IM route may be used

### DSED or VCD

AGE: ≥ 18 years  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: Non-traumatic VF/pulseless VT of presumed cardiac origin  
Three consecutive standard shocks by Paramedics or Fire Services

### amiodarone

AGE: ≥ 24 hours  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: VF **OR** pulseless VT

### lidocaine

AGE: ≥ 24 hours  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: VF **OR** pulseless VT

### 0.9% NaCl Fluid Bolus

AGE: ≥ 24 hours  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: PEA **OR**  
Any other rhythm where hypovolemia is suspected

### Medical TOR

AGE: ≥ 16 years  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: Arrest not witnessed by paramedic **AND** No ROSC after 20 minutes of resuscitation **AND** No defibrillation delivered

## CONTRAINDICATIONS

### CPR

Obviously dead as per  
BLS PCS

Meet conditions of the  
BLS PCS *Do Not  
Resuscitate (DNR)  
Standard*

### Manual Defibrillation

N/A

### AED or SAED Defibrillation

N/A

### EPINEPHrine

Allergy or sensitivity to  
EPINEPHrine

### amiodarone

Allergy or sensitivity to  
amiodarone

### lidocaine

Allergy or sensitivity to  
lidocaine

### 0.9% NaCl Fluid Bolus

Fluid overload

### Medical TOR

Pregnancy presumed to be  $\geq 20$  weeks gestation  
Suspected hypothermia  
Airway obstruction  
Non-opioid drug overdose/toxicity

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **Manual defibrillation**

	Age ≥ 24 hours to < 8 years	Age ≥ 8 years
<i>Dose</i>	1 defibrillation	1 defibrillation
<i>Initial dose</i>	2 J/kg	As per RBHP / manufacturer
<i>Subsequent dose(s)</i>	4 J/kg	As per RBHP / manufacturer
<i>Dosing interval</i>	2 min	2 min
<i>Max. # of doses</i>	N/A	N/A

Consider **AED or SAED defibrillation** (if not using manual defibrillation)

	Age ≥ 24 hours to < 8 years	Age ≥ 8 years
<i>Dose</i>	1 defibrillation with or without pediatric attenuator cable	1 defibrillation
<i>Max. single dose</i>	As per RBHP / manufacturer	As per RBHP / manufacturer
<i>Dosing interval</i>	2 min	2 min
<i>Max. # of doses</i>	N/A	N/A

Consider **DSED** (if authorized) or **VCD** (if DSED is not available or authorized)

	Age ≥ 18 years
<i>Dose</i>	1 DSED or VCD
<i>Max. single dose</i>	As per RBHP / manufacturer
<i>Dosing interval</i>	2 min
<i>Max. # of doses</i>	N/A

Consider **EPINEPHrine** (if anaphylaxis is suspected as the causative event of the cardiac arrest)

Route IM	
Concentration 1 mg/mL = 1:1,000	
<i>Dose</i>	0.01 mg/kg*
<i>Max. single dose</i>	0.5 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

\* The EPINEPHrine dose may be rounded to the nearest 0.05 mg

Consider **EPINEPHrine**

	Age ≥ 24 hours to < 12 years		Age ≥ 12 years	
	Route		Route	
	IV / IO / CVAD	ETT	IV / IO / CVAD	ETT
<i>Solution</i>	0.1 mg/mL = 1:10,000	1 mg/mL = 1:1,000	0.1 mg/mL = 1:10,000	as per RBHP
<i>Dose</i>	0.01 mg/kg*  (0.1 mL/kg)	0.1 mg/kg to a max of 2 mg  (0.1 mL/kg to a max of 2mL)	1 mg	2 mg
<i>Min. single dose</i>	0.05 mg	0.5 mg	1 mg	2 mg
<i>Dosing interval</i>	4 min	4 min	4 min	4 min
<i>Max. # of doses</i>	N/A	N/A	N/A	N/A

\* The EPINEPHrine dose may be rounded to the nearest 0.05 mg

Consider **amiodarone** (if not using lidocaine)

	Age ≥ 24 hours to < 12 years	Age ≥ 12 years
	Route	Route
	IV / IO / CVAD	IV / IO / CVAD
<i>Initial Dose</i>	5 mg/kg	300 mg
<i>Max. initial dose</i>	300 mg	300 mg
<i>Subsequent dose(s)</i>	5 mg/kg	150 mg
<i>Max. repeat dose</i>	150 mg	150 mg
<i>Dosing interval</i>	4 min	4 min
<i>Max. # of doses</i>	2	2

Consider **lidocaine** (if not using amiodarone)

	Age ≥ 24 hours to < 12 years		Age ≥ 12 years	
	Route		Route	
	IV / IO / CVAD	ETT	IV / IO / CVAD	ETT
<i>Initial Dose</i>	1 mg/kg	2 mg/kg	1.5 mg/kg	3 mg/kg
<i>Second Dose</i>	1 mg/kg	2 mg/kg	0.75 mg/kg	1.5 mg/kg
<i>Min. single dose</i>	N/A	N/A	N/A	N/A
<i>Dosing interval</i>	4 min	4 min	4 min	4 min
<i>Max. # of doses</i>	2	2	2	2

Consider **0.9% NaCl fluid bolus**

	Age ≥24 hours to < 12 years	Age ≥ 12 years
	Route	Route
	IV / IO / CVAD	IV / IO / CVAD
<i>Infusion</i>	20 mL/kg	20 mL/kg
<i>Infusion interval</i>	Immediate	Immediate
<i>Reassess every</i>	100 mL	250 mL
<i>Max. volume</i>	2,000 mL	2,000 mL

**Mandatory Provincial Patch Point**

Patch to consider Medical TOR (if applicable).

Patch early to consider TOR if there are extenuating circumstances or where the paramedic considers ongoing resuscitation to be futile.

If the patch fails, and/or, no ROSC after 20 minutes of resuscitation, initiate transport.

**CLINICAL CONSIDERATIONS**

The IV/IO/CVAD routes of medication administration are preferred over the ETT route. However, ETT administration may be used if the IV/IO/CVAD routes are delayed (e.g.  $\geq 5$  min).

The BHP might not authorize TOR even though the patient meets TOR rule. Factors may include: location of the patient, EtCO<sub>2</sub>, age, bystander witnessed, bystander CPR, transportation time, and unusual cause of cardiac arrest such as electrocution, hanging, and toxicology.

DSED/VCD:

The second defibrillator for Dual Sequential Defibrillation will be a paramedic service defibrillator or a fire service defibrillator (in order of preference and if agreed to by the fire service). If a second defibrillator is not available, Vector Change Defibrillation should be provided.



## LOCAL BHP CONSULTATION ADVISORY

In extenuating circumstances during unusual or prolonged codes, Paramedics may choose to patch for consultation. Extenuating circumstances may include, but are not limited to, the following:

1. Unusual cardiac arrest causes (ie. FBAO, hypothermia, electrocution, toxicity)
2. Excessive epinephrine administration (>5-6mg) in prolonged resuscitations.
3. Excessive number of shocks (>3 with vector change) delivered without change in refractory dysrhythmia.

Patient presentation/underlying cause of cardiac arrest should be considered when carrying out a treatment plan.



**NOTE:** Refer to page 35 for **Epinephrine 1mg/mL = 1:1000 IM Dosing Chart**



**NOTE:** Refer to page 61 for **Defibrillation Joule Setting Reference Chart.**



**NOTE:** Refer to page 202 for **CPR Guidelines.**

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Adult Defibrillation Joule Settings Reference

Manufacturer:	Series:	Joule Settings:
<b>Medtronic</b>	Lifepack	200, 300, 360 Joules
<b>Phillips</b>	MRX / FR2	150 Joules non escalating
<b>ZOLL</b>	E, M, or X Series	120, 150, 200 Joules

## Trauma Cardiac Arrest Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Cardiac arrest secondary to severe blunt or penetrating trauma.

### CONDITIONS

<p><b>CPR</b></p> <p>AGE: N/A</p> <p>LOA: Altered</p> <p>HR: N/A</p> <p>RR: N/A</p> <p>SBP: N/A</p> <p>Other: Performed in 2 minute intervals</p>	<p><b>Manual Defibrillation</b></p> <p>AGE: ≥24 hours</p> <p>LOA: Altered</p> <p>HR: N/A</p> <p>RR: N/A</p> <p>SBP: N/A</p> <p>Other: VF <b>OR</b> pulseless VT</p>	<p><b>AED or SAED Defibrillation</b></p> <p>AGE: ≥24 hours</p> <p>LOA: Altered</p> <p>HR: N/A</p> <p>RR: N/A</p> <p>SBP: N/A</p> <p>Other: Defibrillation indicated</p> <p>If not using manual defibrillation</p>
<p><b>Needle thoracostomy</b></p> <p>AGE: N/A</p> <p>LOA: N/A</p> <p>HR: N/A</p> <p>RR: N/A</p> <p>SBP: N/A</p> <p>Other: Suspected tension pneumothorax <b>AND</b> absent or severely diminished breath sound on the affected side(s)</p>	<p><b>Trauma TOR</b></p> <p>AGE: ≥16 years</p> <p>LOA: Altered</p> <p>HR: 0</p> <p>RR: 0</p> <p>SBP: N/A</p> <p>Other: No palpable pulses <b>AND</b> No defibrillation delivered <b>AND</b> Rhythm Asystole <b>AND</b> No signs of life at any time since fully extricated <b>OR</b> Signs of life when fully extricated with the closest ED ≥30 min transport time away <b>OR</b> Rhythm PEA with the closest ED ≥30 min transport time away</p>	

## CONTRAINDICATIONS

### CPR

Obviously dead as per  
BLS PCS

Meet conditions of the  
BLS PCS *Do Not  
Resuscitate (DNR)  
Standard*

### Manual Defibrillation

Rhythms other than VF  
or pulseless VT

### AED or SAED Defibrillation

Non-shockable rhythm

### Needle thoracostomy

N/A

### Trauma TOR

Age <16 years

Defibrillation delivered

Signs of life at any time since fully extricated.

Rhythm PEA and closest ED <30 min transport  
time away

Patients with penetrating trauma to the torso or  
head/neck and Lead Trauma Hospital < 30 min  
transport time away

## TREATMENT

Consider **CPR** as per current Heart and Stroke Foundation of Canada  
Guidelines

Consider **Manual defibrillation** (if available and authorized)

	Age ≥24 hours to <8 years	Age ≥8 years
<i>Dose</i>	1 defibrillation	1 defibrillation
<i>Initial dose</i>	2 J/kg	As per RBHP / manufacturer
<i>Dosing interval</i>	N/A	N/A
<i>Max. # of doses</i>	1	1

Consider **AED or SAED defibrillation** (if not using manual defibrillation)

	Age ≥24 hours to <8 years	Age ≥8 years
<i>Dose</i>	1 defibrillation with or without pediatric attenuator cable	1 defibrillation
<i>Max. single dose</i>	As per RBHP / manufacturer	As per RBHP / manufacturer
<i>Dosing interval</i>	N/A	N/A
<i>Max. # of doses</i>	1	1

Consider **needle thoracostomy**



### Mandatory Provincial Patch Point



Patch to BHP for authorization to apply the Trauma TOR if applicable. If the BHP patch fails, or the Trauma TOR does not apply, transport to the closest appropriate receiving facility following the 1<sup>st</sup> analysis/defibrillation.

## CLINICAL CONSIDERATIONS

If no obvious external signs of significant blunt trauma, consider medical cardiac arrest and treat according to the appropriate medical cardiac arrest directive.

Signs of life: specifically any spontaneous movement, respiratory efforts, organized electrical activity on ECG, and reactive pupils.

An intravenous fluid bolus may be considered, where it does not delay transport and should not be prioritized over management of other reversible pathology.




**NOTE:** Refer to page 61 for **Defibrillation Joule Setting Reference Chart**.



**NOTE:** Refer to page 202 for **CPR Guidelines**.

## Treatment – Algorithm for Trauma Arrest



Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

## Newborn Resuscitation Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Newborn patient.

### CONDITIONS

PPV	CPR	EPINEPHrine
AGE: <24 hours	AGE: <24 hours	AGE: <24 hours
LOA: N/A	LOA: N/A	LOA: N/A
HR: <100 bpm	HR: <60 bpm	HR: <60 bpm
RR: N/A	RR: N/A	RR: N/A
SBP: N/A	SBP: N/A	SBP: N/A
Other: N/A	Other: After 30 seconds of PPV using room air	Other: After 30 seconds of PPV <b>AND</b> 30 seconds of CPR

### CONTRAINDICATIONS

PPV	CPR
Obviously dead as per BLS PCS	Obviously dead as per BLS PCS
Presumed gestational age less than 20 weeks	Presumed gestational age less than 20 weeks
EPINEPHrine	
Allergy or sensitivity to EPINEPHrine	
Presumed gestational age less than 20 weeks	



## TREATMENT

Consider **PPV** as per the treatment flowchart

Consider **CPR** as per current Heart and Stroke Foundation of Canada Guidelines

Consider **EPINEPHrine**

Age

< 24 hours


	Route	
	IV/IO	ETT*
<i>Solution</i>	0.1 mg/mL = 1:10,000	0.1 mg/mL = 1:10,000
<i>Dose</i>	0.01 mg/kg (0.1 mL/kg)	0.1 mg/kg (1.0 mL/kg)
<i>Min. single dose</i>	0.05 mg (0.5 mL)	N/A
<i>Max. single dose</i>	N/A	0.3 mg (3.0 mL)
<i>Dosing interval</i>	4 min	N/A
<i>Max. # of doses</i>	N/A	1

**\* EPINEPHrine is to be administered IV/IO after the single ETT dose if the conditions are still met**

## Treatment

Targeted Preductal SpO<sub>2</sub>  
(Right Hand)

1 min	60% - 65%
2 min	65% - 70%
3 min	70% - 75%
4 min	75% - 80%
5 min	80% - 85%
10 min	85% - 95%



## CLINICAL CONSIDERATIONS

If newborn resuscitation is required, initiate cardiac monitoring and right-hand pulse oximetry monitoring.

Infants born between 20-25 weeks gestation may be stillborn or die quickly. Initiate resuscitation and transport as soon as feasible.

If gestational age cannot be confirmed, initiate resuscitation and rapid transport.

If newborn is less than 20 weeks gestation, resuscitation is futile. Provide the newborn with warmth and consider patching to BHP for further direction.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## APGAR Score Reference

Parameter	0	1	2
<b>Heart rate (bpm)</b>	0 (absent)	Slow (< 100)	$\geq 100$
<b>Respiratory effort</b>	Absent	Slow, irregular	Good, crying
<b>Muscle tone</b>	None, limp	Some flexion	Active motion
<b>Reflex irritability (suction of nares, tactile stimulation)</b>	None	Some grimace	Good grimace, cough, cry
<b>Colour</b>	Blue or pale	Pink body with blue extremities	Completely pink

- ▶ APGAR performed at 1 minute & 5 minutes after delivery
- ▶ Maximum possible total score is 10 (5 parameters x maximum score 2 for each parameter)
- ▶ Don't wait for APGAR to make decision on resuscitation

## Neonatal Pre-ductal Oxygen Saturation Reference

### TARGETED PRE-DUCTAL SpO<sub>2</sub>

#### After Birth

1 min	60-65%
2 min	65-70%
3 min	70-75%
4 min	75-80%
5 min	80-85%
10 min	85-95%

In all neonates, only apply the pulse oximeter to the **RIGHT HAND**.  
Target the above values when:

- ▶ Resuscitation is anticipated
- ▶ PPV is required for more than a few breaths
- ▶ Persistent central cyanosis, or if you need to confirm your perception of central cyanosis
- ▶ Any administration of supplemental oxygen

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain /  
Sed. /  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Return of Spontaneous Circulation (ROSC) Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Patient with return of spontaneous circulation (ROSC) after the resuscitation was initiated.

### CONDITIONS

#### 0.9% NaCl Fluid Bolus

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: Hypotension

Other: Chest auscultation is clear

#### DOPamine

AGE:  $\geq 8$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: Hypotension

Other: N/A

### CONTRAINDICATIONS

#### 0.9% NaCl Fluid Bolus

Fluid overload

#### DOPamine

Allergy or sensitivity to DOPamine

Tachydysrhythmias excluding sinus tachycardia

Mechanical shock

Pheochromocytoma

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **optimizing ventilation and oxygenation**

Titrate oxygenation 94%-98%

Avoid hyperventilation and target  $\text{ETCO}_2$  to 30-40 mmHg with continuous waveform capnography (if available)

Consider **0.9% NaCl fluid bolus**

	Age <12 years	Age ≥ 12 years
	Route	Route
	IV/IO/CVAD	IV/IO/CVAD
<i>Infusion</i>	10 mL/kg	10 mL/kg
<i>Infusion interval</i>	Immediate	Immediate
<i>Reassess every</i>	100 mL	250 mL
<i>Max. volume</i>	1,000 mL	1,000 mL

Consider **DOPamine**

	Age ≥8 years
	Route
	IV
<i>Initial Infusion Rate</i>	5 mcg/kg/min
<i>Titration increment</i>	5 mcg/kg/min
<i>Titration interval</i>	5 min
<i>Max infusion rate</i>	20 mcg/kg/min

**NOTE:** Titrate DOPamine to achieve a SBP of ≥90 to <110mmHg. If discontinuing DOPamine electively, do so gradually over 5-10 minutes.

Consider **12 lead ECG acquisition and interpretation**

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain /  
Sed. /  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## CLINICAL CONSIDERATIONS

Consider initiating transport in parallel with the above treatment.

Adult IO administration of a NaCl bolus requires the ACP to be authorized.

Notify receiving hospital staff if DOPamine drip goes interstitial.



**NOTE:** Refer to page 200 for **12 Lead ECG Placement Reference.**



## Single Strength DOPamine Dosing Chart

DOPamine INFUSION RATE (mL/hr or drops/min with a microdrip set)  
[Using an 800 mcg/mL ('single strength') solution]

Weight (kg)	Drip Rate (drops/min)				
	2 (mcg/kg/minute)	5 (mcg/kg/minute)	10 (mcg/kg/minute)	15 (mcg/kg/minute)	20 (mcg/kg/minute)
5	1	2	4	6	8
10	2	4	8	11	15
15	2	6	11	17	23
20	3	8	15	23	30
25	4	9	19	28	38
30	5	11	23	34	45
35	5	13	26	39	53
40	6	15	30	45	60
45	7	17	34	51	68
50	8	19	38	56	75
55	8	21	41	62	83
60	9	23	45	68	90
65	10	24	49	73	98
70	11	26	53	79	105
75	11	28	56	84	113
80	12	30	60	90	120
85	13	32	64	96	128
90	14	34	68	101	135
95	14	36	71	107	143
100	15	38	75	113	150
105	16	39	79	118	158
110	17	41	83	124	165
115	17	43	86	129	173
120	18	45	90	135	180

## Cardiac Ischemia Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Suspected cardiac ischemia.

### CONDITIONS

ASA	nitroglycerin	morphine
AGE: ≥18 years	AGE: ≥18 years	AGE: ≥18 years
LOA: Unaltered	LOA: Unaltered	LOA: Unaltered
HR: N/A	HR: 60-159 bpm	HR: N/A
RR: N/A	RR: N/A	RR: N/A
SBP: N/A	SBP: Normotension	SBP: Normotension
Other: Able to chew and swallow	Other: Prior history of nitroglycerin use <b>OR</b> IV access obtained	Other: Severe pain

### CONTRAINDICATIONS

ASA	nitroglycerin	morphine
Allergy or sensitivity to NSAIDS	Allergy or sensitivity to nitrates	Allergy or sensitivity to morphine
If asthmatic, no prior use of ASA	Phosphodiesterase inhibitor use within the previous 48 hours	SBP drops by one-third or more of its initial value after morphine is administered
Current active bleeding	SBP drops by one-third or more of its initial value after nitroglycerin is administered	
CVA or TBI in the previous 24 hours	12-lead ECG compatible with Right Ventricular MI	

## TREATMENT


**Patient Drug • Dose Route Time.**

 Consider **ASA**

	Route
	PO
Dose	160-162 mg
Max. single dose	162 mg
Dosing interval	N/A
Max. # of doses	1

 Consider **12-lead ECG acquisition and interpretation for STEMI**

 Consider **nitroglycerin**

	STEMI	
	NO	YES
	SBP	SBP
	≥100 mmHg	≥100 mmHg
	Route	Route
	SL	SL
Dose	0.3 mg <b>OR</b> 0.4 mg	0.3 mg <b>OR</b> 0.4 mg
Max. single dose	0.4 mg	0.4 mg
Dosing interval	5 min	5 min
Max. # of doses	6	3

 Consider **morphine** (after the 3<sup>rd</sup> dose of nitroglycerin or if nitroglycerin is contraindicated)

	Route
	IV
Dose	2 mg
Max. single dose	2 mg
Dosing interval	5 min
Max. # of doses	5

## CLINICAL CONSIDERATIONS

Suspect a Right Ventricular MI in all inferior STEMIs and perform at minimum V4R to confirm (ST-elevation  $\geq 1$ mm in V4R).

Do not administer nitroglycerin to a patient with a Right Ventricular STEMI.

Apply defibrillation pads when a STEMI is identified.

The goal for time to 12-lead ECG from first medical contact is < 10 minutes where possible.



**NOTE:** Refer to page 200 for **12 Lead ECG Placement Reference.**

# Acute Cardiogenic Pulmonary Edema

## Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Moderate to severe respiratory distress;

#### AND

Suspected acute cardiogenic pulmonary edema

### CONDITIONS

#### **nitroglycerin**

AGE:  $\geq 18$  years

LOA: N/A

HR: 60-159 bpm

RR: N/A

SBP: Normotension

Other: N/A

### CONTRAINDICATIONS

#### **nitroglycerin**

Allergy or sensitivity to nitrates

Phosphodiesterase inhibitor use within the previous 48 hours

SBP drops by one-third or more of its initial value after nitroglycerin is administered

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT

**Patient • Drug • Dose • Route • Time.**Consider **nitroglycerin**

	SBP ≥100 mmHg to <140 mmHg		SBP ≥140 mmHg	
	IV or Hx*		IV or Hx*	
	Yes		No	
	Route		Route	
	SL		SL	
<i>Dose</i>	0.3 mg or 0.4 mg		0.3 mg or 0.4 mg	
<i>Max. single dose</i>	0.4 mg		0.4 mg	
<i>Dosing interval</i>	5 min		5 min	
<i>Max. # of doses</i>	6		6	

**\*Hx refers to a patient with a prior history of nitroglycerin use**Consider **12-lead ECG acquisition and interpretation**

## CLINICAL CONSIDERATIONS

N/A

**NOTE: Refer to page 200 for 12 Lead ECG Placement Reference.**

## Cardiogenic Shock Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

STEMI-positive 12-lead ECG;

#### AND

Cardiogenic shock.

### CONDITIONS

#### 0.9% NaCl fluid bolus

AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: Hypotension

Other: Chest auscultation is clear

#### DOPamine

AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: Hypotension

Other: N/A

### CONTRAINDICATIONS

#### 0.9% NaCl fluid bolus

Fluid overload

SBP  $\geq 90$  mmHg

#### DOPamine

Allergy or sensitivity to DOPamine

Tachydysrhythmias excluding sinus tachycardia

Mechanical shock

Hypovolemia

Pheochromocytoma

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **0.9% NaCl fluid bolus**

	Age
	≥18 years
	Route
	IV/IO/CVAD
Infusion	10 mL/kg
Infusion interval	N/A
Reassess every	250 mL
Max. volume	1,000 mL

**NOTE:** If NaCl bolus contraindicated due to pulmonary crackles, consider DOPamine.

Consider **DOPamine**

	Route
	IV
Initial infusion rate	5 mcg/kg/min
Titration increment	5 mcg/kg/min
Titration interval	5 min
Max. infusion rate	20 mcg/kg/min

**NOTE:** Titrate DOPamine to achieve a SBP of ≥90 to <110 mmHg. If discontinuing DOPamine electively, do so gradually over 5-10 minutes.

## CLINICAL CONSIDERATIONS

Contact BHP if patient is bradycardic.



## Symptomatic Bradycardia Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Bradycardia;

**AND**

Hemodynamic instability.

### CONDITIONS

<b>atropine</b>	<b>Transcutaneous Pacing</b>	<b>DOPamine</b>
AGE: $\geq 18$ years	AGE: $\geq 18$ years	AGE: $\geq 18$ years
LOA: N/A	LOA: N/A	LOA: N/A
HR: $< 50$ bpm	HR: $< 50$ bpm	HR: $< 50$ bpm
RR: N/A	RR: N/A	RR: N/A
SBP: Hypotension	SBP: Hypotension	SBP: Hypotension
Other: N/A	Other: N/A	Other: N/A

### CONTRAINDICATIONS

<b>atropine</b>	<b>Transcutaneous Pacing</b>	<b>DOPamine</b>
Allergy or sensitivity to atropine	Hypothermia	Allergy or sensitivity to DOPamine
Hypothermia		Mechanical shock
History of heart transplant		Pheochromocytoma

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **Rhythm determination**

Consider **12 lead ECG acquisition and interpretation** (if this won't delay therapy)

Consider **atropine**

	Route
	IV
<i>Dose</i>	1 mg
<i>Max. single dose</i>	1 mg
<i>Dosing interval</i>	5 min
<i>Max. # of doses</i>	2

Consider **transcutaneous pacing**

Consider **DOPamine**

	Route
	IV
<i>Initial infusion rate</i>	5 mcg/kg/min
<i>Titration increment</i>	5 mcg/kg/min
<i>Titration interval</i>	5 min
<i>Max. infusion rate</i>	20 mcg/kg/min

**NOTE:** Titrate DOPamine to achieve a SBP of  $\geq 90$  to  $< 110$  mmHg. If discontinuing DOPamine electively, do so gradually over 5-10 minutes.

## CLINICAL CONSIDERATIONS

TCP should not be delayed for placement of an IV.

A fluid bolus should be considered with all symptomatic bradycardia patients if indicated.




**NOTE:** Refer to page 76 for **Dopamine Dosing Chart**.



**NOTE:** Refer to page 136 for **Procedural Sedation Medical Directive**

## Transcutaneous Pacing

- ▶ Refer to the Symptomatic Bradycardia Medical Directive for indications, conditions and contraindications for transcutaneous pacing.
- ▶ Ensure the limb leads are applied for cardiac monitoring
- ▶ Place the defibrillation pads on patient as per manufacturer's recommendation. Generally "anterior/posterior" or sternum/apex placements are acceptable.
- ▶ Set the pacing rate to 80 beats per minute.
- ▶ Increase the output or milliamps slowly until electrical capture is confirmed by the presence of pacer spikes followed by a wide QRS complex (see below).



- ▶ Once electrical capture is confirmed, the presence of mechanical capture must be confirmed and consistently monitored.
- ▶ Mechanical capture should be confirmed through palpation of a mechanical pulse that matches the pre-set rate on the pacer.
- ▶ Upon confirmation of mechanical capture, the Paramedic should increase the output or milliamps by 5-10% to ensure mechanical capture is maintained.
- ▶ If pacing does not achieve mechanical capture despite maximal output and good connection between the pads and skin, then this should be discontinued.

### CONSIDERATIONS

- ▶ Generally electrical capture will be achieved between 70 and 120 mA, but higher energy settings will occasionally be required.
- ▶ Many conscious patients who are receiving TCP therapy will require sedation. Consider applying the Procedural Sedation Medical Directive.
- ▶ Ensure continuous monitoring of mechanical capture as it can be lost over time.

## Tachydysrhythmia Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Symptomatic Tachydysrhythmia.

### CONDITIONS

#### Valsalva Maneuver

AGE:  $\geq 18$  years  
LOA: Unaltered  
HR:  $\geq 150$  bpm  
RR: N/A  
SBP: Normotension  
Other: Narrow complex and regular rhythm

#### adenosine

AGE:  $\geq 18$  years  
LOA: Unaltered  
HR:  $\geq 150$  bpm  
RR: N/A  
SBP: Normotension  
Other: Narrow complex and regular rhythm

#### amiodarone

AGE:  $\geq 18$  years  
LOA: Unaltered  
HR:  $\geq 120$  bpm  
RR: N/A  
SBP: Normotension  
Other: Wide complex and regular rhythm

#### lidocaine

AGE:  $\geq 18$  years  
LOA: Unaltered  
HR:  $\geq 120$  bpm  
RR: N/A  
SBP: Normotension  
Other: Wide complex and regular rhythm

#### Synchronized Cardioversion

AGE:  $\geq 18$  years  
LOA: N/A  
HR:  $\geq 120$  bpm (wide) **OR**  
 $\geq 150$  bpm (narrow)  
RR: N/A  
SBP: Hypotension  
Other: Altered mental status, ongoing chest pain, other signs of shock

## CONTRAINDICATIONS

### Valsalva Maneuver

Sinus tachycardia or atrial fibrillation or atrial flutter

### adenosine

Allergy or sensitivity to adenosine

Sinus tachycardia or atrial fibrillation or atrial flutter

Patient taking dipyridamole or carbamazepine

Bronchoconstriction on exam

### amiodarone

Allergy or sensitivity to amiodarone

### lidocaine

Allergy or sensitivity to lidocaine

### Synchronized Cardioversion

N/A

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **rhythm determination** (confirm regularity)

Consider **12-lead ECG acquisition and interpretation to confirm QRS width** (if this won't delay therapy)

Consider **valsalva maneuver**

Perform a maximum of 2 attempts lasting 10 to 20 seconds duration each.

Consider **adenosine**

	Route IV
<i>Initial dose</i>	6 mg
<i>Subsequent dose</i>	12 mg
<i>Dosing interval</i>	2 min
<i>Max. # of doses</i>	2

### **Mandatory Provincial Patch Point**

Patch to BHP for authorization to proceed with amiodarone or lidocaine or if monomorphic wide complex regular rhythm for adenosine.

Consider **amiodarone OR lidocaine** (if not using amiodarone)

	Medication Amiodarone	Medication Lidocaine
	Route IV*	Route IV
<i>Initial dose</i>	150 mg	1.5 mg/kg
<i>Subsequent dose</i>	150 mg	0.75 mg/kg
<i>Max. single dose</i>	150 mg	150 mg
<i>Dosing interval</i>	10 min	10 min
<i>Max. # of doses</i>	2	3

**\*Amiodarone should be administered by IV infusion over 10 min.**

### **Mandatory Provincial Patch Point**

Patch to BHP for authorization to proceed with synchronized cardioversion.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain /  
Sed. /  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Consider **synchronized cardioversion**

Administer up to 3 synchronized shocks in accordance with BHP direction and energy settings. (In the setting of a patch failure, the energy settings to be used are 100 J, 200 J and the maximum manufacturer setting.)

## CLINICAL CONSIDERATIONS

N/A

## CONSIDERATIONS FOR TREAT AND DISCHARGE (IF AUTHORIZED)

The patient must meet all of the following criteria:

- the patient is  $\geq 18$  AND  $< 65$  years old;
- patient must have a prior history of SVT;
- the patient presented with narrow complex and regular rhythm Supraventricular Tachycardia (SVT);
- the patient must have only had a single SVT episode in the past 24 hours
- the patient has returned to normal sinus rhythm (NSR) either spontaneously, with a valsalva maneuver or with adenosine treatment by paramedics and is now asymptomatic;
- the patient has returned to their normal level of consciousness;
- a complete set of vital signs are within expected normal ranges with a HR  $< 100$  bpm and the patient remains in NSR for at least 15 minutes post conversion;

### AND

- the patient was not treated with electrical cardioversion by paramedics;
- the patient is not pregnant;
- the SVT must not be related to alcohol or substance abuse or withdrawal, and;
- the patient has no fever or preceding illness.



In addition to the above criteria, if all of the following requirements have been met, the patient can be discharged by paramedics:

- a responsible adult agrees to remain with the patient for the next 4 hours;
- all of the patient or substitute decision makers questions were answered and a care plan was developed;
- the patient or substitute decision maker has been advised to follow up with their primary health care team or provider;
- clear instructions to call 911 were provided should symptoms redevelop;
- patient or substitute decision maker has the ability to access 911 should symptoms redevelop, and;
- patient or substitute decision maker consents to the discharge.

### CLINICAL CONSIDERATIONS (TREAT AND DISCHARGE)

Patch to BHP for consultation if you are unclear if the patient meets all of the discharge criteria.



*NOTE: Refer to page 136 for **Procedural Sedation Medical Directive***



*NOTE: Refer to page 200 for **12 Lead ECG Placement Reference***

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.


Medic.  
Info.

Contact

Destinat.  
Guide.

## Synchronized Cardioversion

- ▶ Refer to the Tachydysrhythmia Medical Directive for the specific indications, conditions and contraindications.
- ▶ Patch to the BHP is required in accordance with the Medical Directive. In the setting of a patch failure, and the patient otherwise meets the medical directive, the energy settings to be used are 100J, 200J and the maximum manufacturer setting.
- ▶ Ensure the limb leads are applied for cardiac monitoring
- ▶ Place the defibrillation pads on patient as per manufacturer's recommendation. Generally "anterior/posterior" or sternum/apex placements are acceptable.
- ▶ Activate **synchronize** mode on the monitor if necessary.
- ▶ Press the synchronize button and be sure the machine has "marked" each QRS complex. See image below.



- ▶ Once it has been identified that each QRS is appropriately marked, set the energy in accordance with the BHP order, or in the event of a patch failure, in accordance with the Medical Directive.
- ▶ Press the charge button.
- ▶ Once the defibrillator is charged, clear the patient and press and **hold** the shock button until the energy has been delivered.
- ▶ Reassess patient.
- ▶ Generally, the initial joule setting will be 100J. If a successful cardioversion is not achieved at the initial setting, subsequent synchronized cardioversions may be delivered at increased joule settings in accordance with the BHP orders or the Medical Directive. Ensure that the defibrillator is **synched** before any subsequent cardioversions are delivered.

### CONSIDERATIONS

- ▶ Anomalies in the ECG morphology may cause the machine to mistakenly mark non-QRS portions of the ECG. For example peaked T waves may be marked

in the setting of hyperkalemia, or the P and T waves could be marked in low amplitude states. The paramedic must be sure the machine is properly identifying QRS complexes. This is achieved by cycling through all leads looking for the tallest R waves.

- ▶ The defibrillator must be **resynchronized** before every cardioversion attempt.
- ▶ If synchronization is not possible, consult with the BHP during the patch.

## Traumatic Hemorrhage Medical Directive - Auxiliary

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Suspected hemorrhage due to trauma

#### AND

Hemodynamic instability

### CONDITIONS

#### **Tranexamic Acid (TXA)**

AGE:  $\geq 16$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: HR  $\geq 110$  BPM or  
hypotension

## CONTRAINDICATIONS

### Tranexamic Acid (TXA)

Allergy of sensitivity to TXA

Greater than 3 hours from the time of injury to drug administration **OR** unknown time of injury

Isolated head injury

## TREATMENT

### Consider Tranexamic Acid (TXA)

	Route IV	Route IM
<i>Dose</i>	1000 mg	1000 mg
<i>Max. single dose</i>	1000 mg	1000 mg
<i>Dosing interval</i>	N/A	N/A
<i>Max. # of doses</i>	1	1

## CONSIDERATIONS

TXA should not delay transport and should not be prioritized over the management of other reversible causes.

TXA solution for injection should be administered intravenously by slow injection over a period of at least 5 minutes, as rapid administration can cause hypotension.

## Hyperkalemia Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Suspected hyperkalemia in patients at high risk, including:

Currently on dialysis; **OR**

History of end-stage renal disease; **OR**

Relevant incident history (i.e. prolonged crush injury)

### AND

One of the following clinical situations:

Cardiac Arrest; **OR**

Prearrest with 12-lead ECG changes associated with Hyperkalemia.

### CONDITIONS

#### calcium gluconate 10%

AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### salbutamol

AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

## CONTRAINDICATIONS

### calcium gluconate

Allergy of sensitivity to calcium gluconate

### salbutamol

Allergy or sensitivity to salbutamol

Airway /  
Breath.

Cardiac /  
Circula.

## TREATMENT

Consider **12-lead ECG acquisition and interpretation**

Consider **calcium gluconate 10%**

	Route
	IV/IO/CVAD
<i>Dose</i>	1 g (10 mL) over 2-3 minutes
<i>Max. single dose</i>	1g (10 mL)
<i>Dosing interval</i>	5 minutes
<i>Max. # of doses</i>	2 *

**\*Note: an additional 3rd dose may be administered after 30 minutes if the patient improved initially and symptoms meeting the indications recur.**

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Consider **salbutamol**

	Route	
	MDI*	NEB
<i>Dose</i>	1,600 mcg (16 puffs)	10 mg
<i>Max. single dose</i>	1,600 mcg	10 mg
<i>Dosing interval</i>	Immediate	Immediate
<i>Max. # of doses</i>	2	2

**\*1 puff=100mcg**

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Consider **12-lead ECG acquisition and interpretation**

Contact

## CLINICAL CONSIDERATIONS

In the Indications, the pre-arrest patient would present with one or more of the following: hypotension, altered levels of awareness, or symptomatic bradycardia.

12-lead changes suggestive of hyperkalemia are wide and bizarre QRS complexes [ $\geq 120$  ms], peaked T waves, loss of P waves and/or a QRS complex with a "sine wave" appearance. 12-lead acquisition is intended for the patient not in cardiac arrest to establish the QRS duration before and after treatment.

Whenever possible, both calcium gluconate and salbutamol should be administered as the 2 medications have different modes of action.

The action of calcium gluconate is often visible through the normalization of observed ECG changes of hyperkalemia. If ECG changes do not improve, or if they worsen, additional doses may be required. The duration of action is 20-60 minutes: consider repeat dosing if ECG changes recur during extended transport times.

Caution that calcium gluconate should only be administered in an IV/IO/CVAD that is running well.

Calcium gluconate and sodium bicarbonate should not be mixed or administered in the same IV without flushing well.



## Intravenous and Fluid Therapy Medical Directive

An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.

### INDICATIONS

Actual or potential need for intravenous medication **OR** fluid therapy.

### CONDITIONS

IV Cannulation	0.9% NaCl Fluid Bolus
AGE: N/A	AGE: N/A
LOA: N/A	LOA: N/A
HR: N/A	HR: N/A
RR: N/A	RR: N/A
SBP: N/A	SBP: Hypotension
Other: N/A	Other: N/A

### CONTRAINDICATIONS

IV Cannulation	0.9% NaCl Fluid Bolus
Suspected fracture proximal to the access site	Fluid overload

### TREATMENT

Consider **IV cannulation**

Consider **0.9% NaCl** maintenance infusion

	Age <12 years	Age ≥12 years
	Route	Route
	IV / IO / CVAD	IV / IO / CVAD
<i>Infusion</i>	15 mL/hr	30-60 mL/hr
<i>Infusion interval</i>	N/A	N/A
<i>Reassess every</i>	N/A	N/A
<i>Max. volume</i>	N/A	N/A

**Mandatory Provincial Patch Point**

Patch to BHP for authorization to administer 0.9% NaCl bolus to hypotensive patients <12 years with suspected Diabetic Ketoacidosis (DKA).

**Consider 0.9% NaCl fluid bolus**

	Age <12 years	Age ≥12 years
	Route	Route
	IV / IO / CVAD	IV / IO / CVAD
<i>Infusion</i>	20 mL/kg	20 mL/kg
<i>Infusion interval</i>	Immediate	Immediate
<i>Reassess every</i>	100 mL	250 mL
<i>Max. volume*</i>	2,000 mL	2,000 mL

**\*The maximum volume of 0.9% NaCl is lower for patients in cardiogenic shock and return of spontaneous circulation.**

**CLINICAL CONSIDERATIONS**

Adult IO and CVAD procedures are auxiliary Medical Directives described elsewhere. Fluid administration via the IO or CVAD routes only apply to paramedics authorized to perform these procedures.

Microdrips and/or volume control administration sets should be considered when IV/CVAD access is indicated for patients <12 years of age.

An intravenous fluid bolus may be considered for a patient who does not meet trauma TOR criteria, where it does not delay transport and should not be prioritized over management of other reversible causes.

## Adult Intraosseous Medical Directive –AUXILIARY

*An Advanced Care Paramedic may provide the treatment prescribed in this auxiliary Medical Directive if authorized.*

### INDICATIONS

Actual or potential need for intravenous medication **OR** fluid therapy;  
**AND**  
 IV access is unobtainable;  
**AND**  
 Cardiac arrest or pre-arrest state.

### CONDITIONS

#### IO

AGE: ≥12 years  
 LOA: N/A  
 HR: N/A  
 RR: N/A  
 SBP: N/A  
 Other: N/A

### CONTRAINDICATION

#### IO

Fracture or crush injuries proximal to the access site.  
 Suspected or known replacement / prostheses immediately proximal to the access site

### TREATMENT

Consider **IO access**

### CLINICAL CONSIDERATIONS

N/A



**NOTE: Refer to page 105 to 108 for Intraosseous Site Guidelines**

## Pediatric Intraosseous Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Actual or potential need for intravenous medication **OR** fluid therapy;

**AND**

Intravenous access is unobtainable;

**AND**

Cardiac arrest or pre-arrest state.

### CONDITIONS

#### IO

AGE: <12 years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATION

#### IO

Fracture or crush injuries proximal  
to the access site

Suspected or known replacement  
/ prosthesis proximal to the  
access site.

## TREATMENT

Consider **IO access**

## CLINICAL CONSIDERATIONS

N/A



*NOTE: Refer to page 105 to 108 for **Intraosseous Site Guidelines***

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.


Contact

Destinat.  
Guide.

## INTRAOSSUEOUS SITE GUIDELINES

### PROXIMAL TIBIAL SITE (ADULT AND PEDIATRIC)

- ▶ Refer to the Adult and Pediatric Intraosseous Medical Directives for indications, conditions and contraindications.
- ▶ To landmark this site, rotate the leg externally, then find the medial and lateral condyles of the proximal tibia. Between the condyles, on the top of the anterior tibial crest is the tibial tuberosity. The insertion site is one to two fingerbreadths below and medial to the tibial tuberosity. The needle should be aimed slightly distally away from the growth plate.



- ▶ Cleanse the needle insertion site with aseptic technique.
- ▶ Select the appropriate size IO needle as per the age of the patient and/or the manufacturers recommendation.
- ▶ Insert the IO needle as per the manufacturer's recommendations. For a manual device, place the needle at 90 degrees to the landmark and applying a firm pressure with a screwing-type motion. A distinct tactile "popping" should be felt as needle passes through the bone into the medullary cavity.
- ▶ Once the IO needle has been placed into the medullary cavity of the bone, place the introducer needle into the biohazard container.
- ▶ Confirm proper placement by confirming patency of the IO site, and ensure there are no signs of infiltration. Generally, the posterior aspect of the limb will best show the most prominent signs of infiltration.

- ▶ Attach a primed infusion set to the IO device and set the appropriate infusion rate.
- ▶ Appropriately pad and secure the IO site to maintain stability during patient movement

## CONSIDERATIONS

- ▶ The solution set attached to an IO may require additional pressure to initiate and maintain flow. The paramedic may choose to use a pressure bag, or infuse a predetermined amount of fluid via syringe. When a pressure bag is selected, the flow rate must be constantly monitored as the rate can tend to increase as the compartment becomes more accepting to fluid, and slow as the pressure in the bag decreases.
- ▶ If the first IO attempt is unsuccessful, consider an attempt at the proximal tibial site on the other leg.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure


Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## PROXIMAL HUMERUS SITE IDENTIFICATION – ADULT



- ▶ Select and insert the appropriate size IO needle as per the patient and/or the manufacturers recommendation.

## PROXIMAL HUMERUS NEEDLE SELECTION

- ▶ Palpate and identify which site you want to use. Use the proximal tibia site for pediatrics, and either the proximal tibia or humeral site for adults.
- ▶ Although weight of the patient is one of the criteria for needle selection, the correct needle set is based primarily on the site selected and tissue depth overlying the bone
- ▶ The 45mm needle is recommended for humeral site insertion
- ▶ Due to the anterolateral location of the proximal humerus – use a 45° angle for the insertion
- ▶ Needle Insertion
  - ▶ Stabilize the bone and surrounding tissue (two person technique is best)
  - ▶ Penetrate the skin through to the bone **WITHOUT** running the driver
  - ▶ Assure the 5mm mark is visible when the needle tip has come in contact with the bone
  - ▶ If the 5mm mark is not visible **DO NOT** proceed, the needle is **NOT** long enough!
  - ▶ Run the driver constantly with mild pressure for adult patients, no pressure with pediatric patients



## POST-NEEDLE INSERTION

- ▶ Unscrew the needle (counter clockwise) from the hub and remove it using two hands
- ▶ Connect using the EZ connect adapter ensuring the tubing as been primed with normal saline. DO NOT attach directly to a syringe.
- ▶ Aspirate a small amount of blood (not always possible, does not rule out proper placement) and assess stability of needle in bone to confirm placement
- ▶ Flush with 3-10mL normal saline. In neonates will only need 3mL. In adults use 5-10mL.
- ▶ A second flush may be needed on larger patients
- ▶ Check for any leakage or extravasation or fluid gathering in extremity compartments
- ▶ Put the arm band on the patient's wrist

### NOTE ON FLUSH:

- **NO FLUSH = NO FLOW.**

*Failure to properly flush the IO catheter may result in limited or no flow*

### NOTE ON PRESSURE INFUSER:

- *Pressure in the IV bag must be higher than the pressure inside the bone to achieve flow*
- *Therefore, fluids or medications must be delivered under pressure to obtain maximum flow rates*

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank

# Level of Consciousness

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



## Hypoglycemia Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Suspected hypoglycemia

### CONDITIONS

#### dextrose

AGE: N/A  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: Hypoglycemia

#### glucagon

AGE: N/A  
(≥4 years for IN powder)  
LOA: Altered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: Hypoglycemia

### CONTRAINDICATIONS

#### dextrose

Allergy or sensitivity to dextrose

#### glucagon

Allergy or sensitivity to glucagon  
Pheochromocytoma

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **glucometry**

Consider **dextrose** (D50W diluted as required if not using D10W)

	Age	Age	
	< 2 years	≥ 2 years	
	Concentration	Concentration	Concentration
	10% dextrose	10% dextrose	50% dextrose
	Route	Route	Route
	IV	IV	IV
<i>Dose</i>	0.2 g/kg (2 ml/kg)	0.2 g/kg (2ml/kg)	0.5 g/kg (1 ml/kg)
<i>Max. single dose</i>	5 g (50 ml)	25 g (250 ml)	25 g (50 ml)
<i>Dosing interval</i>	10 min	10 min	10 min
<i>Max. # of doses</i>	2	2	2

**\*Titrate dextrose to a level of awareness where the patient can safely consume complex carbohydrate.**

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Consider **glucagon** (if not using dextrose)

intranasal powder

(if authorized and  
available)

	Age N/A		Age ≥ 4 years
	Weight < 25 kg	Weight ≥ 25 kg	Weight N/A
	Route	Route	Route
	IM	IM	IN
<i>Dose</i>	0.5 mg	1 mg	3 mg
<i>Max. single dose</i>	0.5 mg	1 mg	3 mg
<i>Dosing interval</i>	20 min	20 min	20 min
<i>Max. # of doses</i>	2	2	2

## CLINICAL CONSIDERATIONS

If the patient responds to dextrose or glucagon, he/she may receive oral glucose or other simple carbohydrates.

If only mild signs or symptoms are exhibited, the patient may receive oral glucose or other simple carbohydrates instead of dextrose or glucagon.

If a patient initiates an informed refusal of transport, a final set of vital signs including blood glucometry must be attempted and documented.

Intranasal glucagon is a powder that is supplied in a commercially available single-dose intranasal device.

## CONSIDERATIONS FOR TREAT AND DISCHARGE (IF AUTHORIZED)

All of the following criteria must be met:

- the patient is ≥18 AND <65 years old;
- the patient has a diagnosis of diabetes;
- the hypoglycemia can be explained by insulin administration with inadequate oral intake;

- the hypoglycemia promptly responded to a single administration of dextrose or glucagon as per the Medical Directive and/or consumed oral glucose or other complex carbohydrates; this was a single isolated episode of symptomatic hypoglycemia within the past 24 hours;
- the blood glucose is  $\geq 4.0$  mmol/L after treatment;
- the patient has returned to their normal level of consciousness and is asymptomatic;
- a complete set of vital signs are within expected normal ranges;

#### AND

- not an intentional overdose;
- the hypoglycemia must not be related to alcohol or substance abuse or withdrawal;
- no seizure or reported history of seizure prior to paramedic treatment;
- not on an oral hypoglycemic medication;
- hypoglycemia is not considered to be related to an acute medical illness, and;
- the patient is not pregnant.

In addition to the above criteria, if all of the following requirements have been met, the patient can be discharged by paramedics:

- the patient has access to appropriate carbohydrates;
- a responsible adult agrees to remain with the patient for the next 4 hours;
- all of the patient or substitute decision maker's questions were answered and a care plan was developed;
- the patient or substitute decision maker has been advised to follow up with their primary health care team or provider;
- clear instructions to call 911 were provided should symptoms redevelop;
- patient or substitute decision maker has the ability to access 911 should symptoms redevelop, and
- patient or substitute decision maker consents to the discharge.

### CLINICAL CONSIDERATIONS (TREAT AND DISCHARGE)

Patch to BHP for consultation if you are unclear if the patient meets all of the discharge criteria.

## Dextrose Dosing Guide

Age	Weight kg	Blood Sugar mmol/L	Dextrose prep	Initial dose / Repeat dose		
				Dose g/kg	Volume mL/kg	Amt mL
< 30 days	2	< 3.0	<b>D10W</b> Waste 40 mLs replace w/ sterile water	0.2	2	4
	3				2	6
	4				2	8
	5				2	10
≥30 days to < 2 years	3	< 3.0	<b>D25W</b> Waste 25 mLs replace w/ sterile water	0.5	2	6
	4				2	8
	5				2	10
	6				2	12
	8				2	16
	10				2	20
	12				2	24
	14				2	28
≥ 2 years	10	< 4.0	<b>D50W</b>	0.5	1	10
	15				1	15
	20				1	20
	25				1	25
	30				1	30
	35				1	35
	40				1	40
	45				1	45
	> 50				1	50

5Rs

Patient • Drug • Dose • Route • Time.



## Seizure Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Active generalized motor seizure.

### CONDITIONS

#### midazolam

AGE: N/A

LOA: Unresponsive

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### midazolam

Allergy or sensitivity to midazolam

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **midazolam**

	Route	
	IV / IO	IM / IN / Buccal
<i>Dose</i>	0.1 mg/kg	0.2 mg/kg
<i>Max. single dose</i>	5 mg	10 mg
<i>Dosing interval</i>	5 min	5 min
<i>Max. # of doses</i>	2	2

## CLINICAL CONSIDERATIONS

Conditions such as cardiac arrest and hypoglycemia often present as seizure and should be considered by a paramedic.

Do not delay midazolam administration for blood glucometry in cases where hypoglycemia is not thought to be the causative agent.

Blood glucose should be routinely checked in patients who do not respond to midazolam or have not returned to their baseline LOA after a seizure.

## CONSIDERATIONS FOR TREAT AND DISCHARGE (IF AUTHORIZED)

All of the following criteria must be met:

- the patient is  $\geq 18$  AND  $< 65$  years old;
- patient must have a history of epilepsy;
- the patient is taking their anticonvulsant medication as prescribed;
- the patient must have only had a single seizure episode in the past 24 hours;
- the seizure pattern and duration must be similar to past seizures;
- the patient has returned to their normal level of consciousness and is asymptomatic;
- a complete set of vital signs including temperature are within expected normal ranges;

**AND**

- the seizure must not be related to hypoglycemia, alcohol or substance abuse or withdrawal;
- the patient must not have received midazolam by paramedics;
- the patient did not injure themselves during seizure activity;
- the patient must not have a fever, preceding illness or recently started a new medication, and;
- the patient is not pregnant.

In addition to the above criteria, if all of the following requirements have been met, the patient can be discharged by paramedics:

- a responsible adult agrees to remain with the patient for the next 4 hours;
- all of the patient or substitute decision makers questions were answered and a care plan was developed;
- the patient or substitute decision maker has been advised to follow up with their primary health care team or provider;
- clear instructions to call 911 were provided should symptoms redevelop;
- patient or substitute decision maker has the ability to access 911 should symptoms redevelop, and;
- patient or substitute decision maker consents to the discharge.

## CLINICAL CONSIDERATIONS (TREAT AND DISCHARGE)

Patch to BHP for consultation if you are unclear if the patient meets all of the discharge criteria.

# Seizure Medical Directive Dosing Guide

## Midazolam Dosing Guide

Age	Weight	Route: IM/IN/Buccal			Route: IV/IO		
		Dose: 0.2 mg/kg Supplied: 10 mg/2 mL Use 1 mL syringe Undiluted			Dose: 0.1 mg/kg Supplied: 10 mg/2 mL Use 10 mL syringe diluted to 1 mg/mL		
		Dose	Calculated Volume	Volume to Administer (rounded)	Dose	Actual Volume	Volume to Administer (rounded)
Neonate	3 kg	0.6 mg	0.12 mL	0.10 mL	0.3 mg	0.3 mL	0.4 mL
< 1	6 kg	1.2 mg	0.24 mL	0.25 mL	0.6 mg	0.6 mL	0.6 mL
1	12 kg	2.4 mg	0.48 mL	0.50 mL	1.2 mg	1.2 mL	1.2 mL
2	14 kg	2.8 mg	0.56 mL	0.55 mL	1.4 mg	1.4 mL	1.4 mL
3	16 kg	3.2 mg	0.64 mL	0.65 mL	1.6 mg	1.6 mL	1.6 mL
4	18 kg	3.6 mg	0.72 mL	0.70 mL	1.8 mg	1.8 mL	1.8 mL
5	20 kg	4.0 mg	0.80 mL	0.80 mL	2.0 mg	2.0 mL	2.0 mL
6	22 kg	4.4 mg	0.88 mL	0.90 mL	2.2 mg	2.2 mL	2.2 mL
		Supplied: 10 mg/2 mL Use 3 mL or 10 mL syringe Undiluted			Supplied: 10 mg/2 mL Use 10 mL syringe Diluted to 1 mg/mL		
7	24 kg	4.8 mg	0.96 mL	1.0 mL	2.4 mg	2.4 mL	2.4 mL
8	26 kg	5.2 mg	1.04 mL	1.0 mL	2.6 mg	2.6 mL	2.6 mL
9	28 kg	5.6 mg	1.12 mL	1.2 mL	2.8 mg	2.8 mL	2.8 mL
10	30 kg	6 mg	1.20 mL	1.2 mL	3.0 mg	3.0 mL	3.0 mL
11	32 kg	6.4 mg	1.28 mL	1.2 mL	3.2 mg	3.2 mL	3.2 mL
12	34 kg	6.8 mg	1.36 mL	1.4 mL	3.4 mg	3.4 mL	3.4 mL
	40 kg	8 mg	1.60 mL	1.6 mL	4.0 mg	4.0 mL	4.0 mL
	45 kg	9 mg	1.80 mL	1.8 mL	4.5 mg	4.5 mL	4.5 mL
Max	>50 kg	10 mg	2.00 mL	2.0 mL	5.0 mg	5.0 mL	5.0 mL

Note: Dosage administered can be calculated by the weight based calculation in the Medical Directive and/or by using the above chart. Administered dosage in the chart may be rounded to the nearest volume increment that can be accurately measured.

**Note:**

Dosing for Adult Procedural Sedation: up to 0.1mg/kg (IV/IM/IN); max single dose 5mg; max 2 doses

Dosing for Adult Combative Patient up to 0.1mg/kg (IV/IO/CVAD/IN); max single dose 5mg; max total dose 10mg

Level of Consciousness (LOC) Seizure Medical Directive Dosing Guide v3

# OpioiD Toxicity and Withdrawal

## Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Suspected opioiD toxicity.

### CONDITIONS

#### **naloxone**

AGE:  $\geq 24$  hours

LOA: Altered

HR: N/A

RR:  $<10$  breaths/min

SBP: N/A

Other: Inability to adequately ventilate **OR** persistent need to assist ventilations

#### **buprenorphine/naloxone**

AGE:  $\geq 16$

LOA: Unaltered

HR: N/A

RR: N/A

SBP: N/A

Other: Received naloxone for current opioiD toxicity episode  
**AND**  
Patient is exhibiting acute withdrawal with a COWS\* score  $\geq 8$

### CONTRAINDICATIONS

#### **naloxone**

Allergy or sensitivity to naloxone

#### **buprenorphine/naloxone**

Allergy or sensitivity to buprenorphine

Taken methadone in the past 72 hours

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT

### Consider **naloxone**

	Route IV / IO	Route IM	Route IN	Route SC
<i>Dose</i>	Up to 0.4 mg**	0.4 mg	2-4 mg	0.8 mg
<i>Max. single dose</i>	0.4 mg	0.4 mg	2-4 mg	0.8 mg
<i>Dosing interval</i>	5 mins	5 min	5 min	5 min
<i>Max. # of doses</i>	3	3	3	3

**\*\*For the IV route, titrate naloxone only to restore the patient's respiratory status.**

### Consider **buprenorphine/naloxone** (if available and authorized)

	Route BUC/SL
<i>Initial dose</i>	16 mg
<i>Subsequent dose(s) dose</i>	8 mg
<i>Dosing interval</i>	10 minutes
<i>Max. cumulative dose</i>	24 mgs

## CLINICAL CONSIDERATIONS

Upfront aggressive management of the airway is paramount and the initial priority.

If no response to initial treatment; consider patching for further doses.

If the patient does not respond to airway management and the administration of naloxone, glucometry should be considered.

Combative behaviour should be anticipated following naloxone administration and paramedics should protect themselves accordingly, thus the importance of gradual titrating (if given IV) to desired clinical effect: respiratory rate  $\geq 10$ , adequate airway and ventilation, not full alertness

## \*Clinical Opiate Withdrawal Scale (COWS)

< 5 – No active withdrawal	13–24 – Moderate withdrawal	> 36 – Severe withdrawal
5–12 – Mild withdrawal	25–36 – Moderately severe withdrawal	

A score of  $\geq 8$  is an indication for buprenorphine/naloxone administration

<b>Resting Pulse Rate</b> _____ beats/minute <i>Measured after patient is sitting or lying for one minute</i> 0 pulse rate 80 or below 1 pulse rate 81–100 2 pulse rate 101–120 4 pulse rate greater than 120	<b>GI Upset over last ½ hour</b> 0 no GI symptoms 1 stomach cramps 2 nausea or loose stool 3 vomiting or diarrhea 5 multiple episodes of diarrhea or vomiting
<b>Sweating over past ½ hour not accounted for by room temperature or patient activity</b> 0 no report of chills or flushing 1 subjective report of chills or flushing 2 flushed or observable moistness on face 3 beads of sweat on brow or face 4 sweat streaming off face	<b>Tremor observation of outstretched hands</b> 0 no tremor 1 tremor can be felt, but not observed 2 slight tremor observable 4 gross tremor or muscle twitching
<b>Restlessness observation during assessment</b> 0 able to sit still 1 reports difficulty sitting still, but is able to do so 3 frequent shifting or extraneous movements of legs/arms 5 unable to sit still for more than a few seconds	<b>Yawning observation during assessment</b> 0 no yawning 1 yawning once or twice during assessment 2 yawning three or more times during assessment 4 yawning several times/minute
<b>Pupil Size</b> 0 pupils pinned or normal size for room light 1 pupils possibly larger than normal for room light 2 pupils moderately dilated 5 pupils so dilated that only the rim of the iris is visible	<b>Anxiety or Irritability</b> 0 none 1 patient reports increasing irritability or anxiousness 2 patient obviously irritable anxious 4 patient so irritable or anxious that participation in the assessment is difficult
<b>Bone or Joint Aches</b> <i>If patient was having pain previously, only the additional component attributed to opiates withdrawal is scored</i> 0 not present 1 mild diffuse discomfort 2 patient reports severe diffuse aching of joints/muscles 4 patient is rubbing joints or muscles and is unable to sit still because of discomfort	<b>Gooseflesh Skin</b> 0 skin is smooth 3 piloerection of skin can be felt or hairs standing up on arms 5 prominent piloerection
<b>Runny Nose or Tearing</b> <i>Not accounted for by cold symptoms or allergies</i> 0 not present 1 nasal stuffiness or unusually moist eyes 2 nose running or tearing 4 nose constantly running or tears streaming down cheeks	<div style="text-align: right;"><b>Total Score</b> _____</div> <p><i>The total score is the sum of all 11 items.</i></p> <p>Initials of person completing assessment: _____</p>

## Suspected Adrenal Crisis Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

A patient with primary adrenal failure who is experiencing clinical signs of adrenal crisis.

### CONDITIONS

#### hydrocortisone

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Paramedics are presented with a vial of hydrocortisone for the identified patient **AND**

Age-related hypoglycemia **OR**

GI symptoms (vomiting, diarrhea, abdominal pain) **OR**

Syncope **OR**

Temperature  $\geq 38^{\circ}\text{C}$  or suspected/history of fever **OR**

Altered level of awareness **OR**

Age-related tachycardia **OR**

Age related hypotension

### CONTRAINDICATIONS

#### hydrocortisone

Allergy or sensitivity to  
hydrocortisone



## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **hydrocortisone**

	Route
	IM/IV/IO/CVAD
<i>Dose</i>	2 mg/kg
<i>Max. single dose</i>	100 mg
<i>Dosing interval</i>	N/A
<i>Max. # doses</i>	1

**\*Dose should be rounded to the nearest 10 mg**

## CLINICAL CONSIDERATIONS

N/A

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank

# Pain/Sedation/Nausea

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



## Analgesia Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Pain

### CONDITIONS

#### acetaminophen

AGE: ≥ 12 years

LOA: Unaltered

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### ibuprofen

AGE: ≥ 12 years

LOA: Unaltered

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### ketorolac

AGE: ≥ 12 years

LOA: Unaltered

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### morphine

AGE: ≥ 1 year

LOA: Unaltered

HR: N/A

RR: N/A

SBP: Normotension

Other: N/A

<b>fentaNYL</b>	<b>ketamine</b>
AGE: ≥1 years	AGE: ≥1 years
LOA: Unaltered	LOA: Unaltered
HR: N/A	HR: N/A
RR: N/A	RR: N/A
SBP: N/A	SBP: N/A
Other: N/A	Other: N/A

## CONTRAINDICATIONS

<b>acetaminophen</b>	<b>ibuprofen</b>
Acetaminophen use within previous 4 hours	NSAID use within previous 6 hours
Allergy or sensitivity to acetaminophen	Allergy or sensitivity to ASA or NSAIDs
Hx of liver disease	Patient on anticoagulation therapy
Active vomiting	Current active bleeding
Unable to tolerate oral medication	Hx of peptic ulcer disease or GI bleed
Suspected ischemic chest pain	Pregnant
	If asthmatic, no prior use of ASA or other NSAIDs
	CVA or TBI in the previous 24 hours
	Known renal impairment
	Active vomiting
	Unable to tolerate oral medication
	Suspected ischemic chest pain

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.**ketorolac**

NSAID use within previous 6 hours  
 Allergy or sensitivity to ASA or NSAIDs  
 Patient on anticoagulation therapy  
 Current active bleeding  
 Hx of peptic ulcer disease or GI bleed  
 Pregnant  
 If asthmatic, no prior use of ASA or other NSAIDs  
 CVA or TBI in the previous 24 hours  
 Known renal impairment  
 Suspected ischemic chest pain

**morphine**

Allergy or sensitivity to morphine  
 Treatment of headache  
 Treatment of chronic pain  
 SBP drops by one-third or more of its initial value after morphine is administered  
 Suspected ischemic chest pain (refer to Cardiac Ischemia Medical Directive for suspected cardiac ischemia)  
 Active labour

**fentaNYL**

Allergy or sensitivity to fentaNYL  
 Treatment of headache  
 Treatment of chronic pain  
 SBP drops by one-third or more of its initial value after fentaNYL is administered  
 Suspected ischemic chest pain  
 Active labour

**ketamine**

Allergy or sensitivity to ketamine  
 Treatment of headache  
 Treatment of chronic pain  
 Suspected ischemic chest pain  
 Active labour

**TREATMENT**Consider **acetaminophen**

	Age ≥ 12 years to < 18 years	Age ≥ 18 years
Route	PO	PO
Dose	500-650 mg	960-1,000 mg
Max. single dose	650 mg	1,000 mg
Dosing interval	N/A	N/A
Max. # doses	1	1

Consider **ibuprofen**

Route	Age ≥ 12 years
	PO
Dose	400 mg
Max. single dose	400 mg
Dosing interval	N/A
Max. # doses	1

Consider **ketorolac**

Route	Age ≥ 12 years
	IM / IV
Dose	10-15 mg
Max. single dose	15 mg
Dosing interval	N/A
Max. # doses	1

**⚠ Mandatory Provincial Patch Point ⚠**

Patch to BHP for authorization and dosage verification before:

Administering morphine or fentaNYL for children < 12 years old.

Administering ketamine to patients < 18 years of age.

Consider **fentaNYL (if available and authorized)**

Route	Age ≥1 year to <18 years	Age ≥ 18 years
	IV / IN	IV / IN
Dose	Up to 1 mcg/kg	25 – 75 mcg
Max. single dose	75 mcg	75 mcg
Dosing interval	5 min	5 mins
Max. # of doses	N/A	N/A
Max. cumulative dose	200 mcg	200 mcg

Consider **morphine**

Route	Age ≥1 year to <18 years	Age ≥ 18 years
	IV / SC	IV / SC
Dose	0.05-0.1 mg/kg	2 - 10 mg
Max. single dose	5 mg	10 mg
Dosing interval	15 min	15 min
Max. # of doses	N/A	N/A
Max. cumulative dose	10 mg	20 mg

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Consider **ketamine**

Route	Age ≥1 year to < 18 years		Age ≥ 18 years	
	IV	IN	IV	IN
<i>Dose</i>	0.25 mg/kg	1 mg/kg	0.25 mg/kg	1 mg/kg
<i>Max. single dose</i>	10 mg	30 mg	20 mg	75 mg
<i>Dosing interval</i>	15 min	15 min	15 min	15 min
<i>Max. # of doses</i>	2	2	2	2

## CLINICAL CONSIDERATIONS


Administration of morphine or fentaNYL and ketamine must be sequential, not co-administered. The dosing interval must be no earlier than the most recently administered medication dosing interval.

When higher doses of morphine (5-10 mg) or fentaNYL (50-75 mcg) are given intravenously, consider administering medication in small aliquots q 3 minutes until desired effect or max. single dose is reached to avoid nausea and vomiting.



# Pain Scale Reference

Can be utilized for patients 3 years of age and older.



Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure


Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

# Analgesia Medical Directive - Adult & Pediatric

Age	Weight	Route: Subcutaneous Pediatric dosage 0.05 mg/kg Supplied: 10 mg/mL Use 1 mL Syringe Undiluted			Route: Intravenous Pediatric dosage 0.05 mg/kg Supplied: 10 mg/mL Use 1 mL Syringe Diluted to 1 mg/mL		
		Dose	Calculated Volume	Volume To Administer (rounded)	Dose	Calculated Volume	Volume To Administer (rounded)
		<div><div></div><div>Mandatory Provincial Patch Point</div></div>					
		For patients < 12 years					
Neonate	3 kg	0.15 mg	0.015 mL	---- mL	0.15 mg	0.15 mL	0.15 mL
<1	6 kg	0.3 mg	0.03 mL	0.05 mL	0.3 mg	0.3 mL	0.3 mL
1	12 kg	0.6 mg	0.06 mL	0.05 mL	0.6 mg	0.6 mL	0.6 mL
2	14 kg	0.7 mg	0.07 mL	0.05 mL	0.7 mg	0.7 mL	0.7 mL
3	16 kg	0.8 mg	0.08 mL	0.10 mL	0.8 mg	0.8 mL	0.8 mL
4	18 kg	0.9 mg	0.09 mL	0.10 mL	0.9 mg	0.9 mL	0.9 mL
5	20 kg	1.0 mg	0.10 mL	0.10 mL	1.0 mg	1.0 mL	1.0 mL
6	22 kg	1.1 mg	0.11 mL	0.10 mL	1.1 mg	1.1 mL	1.0 mL
7	24 kg	1.2 mg	0.12 mL	0.1 mL	1.2 mg	1.2 mL	1.2 mL
8	26 kg	1.3 mg	0.13 mL	0.1 mL	1.3 mg	1.3 mL	1.4 mL
9	28 kg	1.4 mg	0.14 mL	0.1 mL	1.4 mg	1.4 mL	1.4 mL
10	30 kg	1.5 mg	0.15 mL	0.2 mL	1.5 mg	1.5 mL	1.6 mL
11	32 kg	1.6 mg	0.16 mL	0.2 mL	1.6 mg	1.6 mL	1.6 mL
		Supplied: 10 mg/mL Use 1 mL Syringe Undiluted			Supplied: 10 mg/mL Use 10 mL Syringe Diluted to 1 mg/mL		
Youth (12-17)	34 kg	1.7 mg	0.17 mL	0.2 mL	1.7 mg	1.7 mL	1.8 mL
	40 kg	2.0 mg	0.20 mL	0.2 mL	2.0 mg	2.0 mL	2.0 mL
	45 kg	2.25 mg	0.225 mL	0.2 mL	2.25 mg	2.25 mL	2.2 mL
	50 kg	2.5 mg	0.25 mL	0.3 mL	2.5 mg	2.5 mL	2.6 mL
	55 kg	2.75 mg	0.275 mL	0.3 mL	2.75 mg	2.75 mL	2.8 mL
	60 kg	3.0 mg	0.30 mL	0.3 mL	3.0 mg	3.0 mL	3.0 mL
	65 kg	3.25 mg	0.325 mL	0.3 mL	3.25 mg	3.25 mL	3.2 mL
	70 kg	3.5 mg	0.35 mL	0.4 mL	3.5 mg	3.5 mL	3.6 mL
	75 kg	3.75 mg	0.375 mL	0.4 mL	3.75 mg	3.75 mL	3.8 mL
	80 kg	4.0 mg	0.40 mL	0.4 mL	4.0 mg	4.0 mL	4.0 mL
	85 Kg	4.25 mg	0.425 mL	0.4 mL	4.25 mg	4.25 mL	4.2 mL
	90 kg	4.5 mg	0.45 mL	0.5 mL	4.5 mg	4.5 mL	4.6 mL
	95 kg	4.75 mg	0.475 mL	0.5 mL	4.75 mg	4.75 mL	4.8 mL
Pediatric Maximum Single Dose		5 mg	0.50 mL	0.5 mL	5.0 mg	5 mL	5 mL

Dosing Interval: **15 minutes** Pediatric Max # of Doses: **4**

Pain/Sedation/Nausea Analgesia Medical Directive - Morphine Dosing Guide v3

# Analgesia Medical Directive - Adult & Pediatric

Dosing Interval: **15 minutes** Pediatric **Max # of Doses: 4**

		Supplied: 10 mg/mL Use 1 mL Syringe Undiluted		Supplied: 10 mg/mL Use 10 mL Syringe Diluted to 1 mg/mL	
<b>Adult</b>	<b>N/A</b>	<b>2 - 10mg</b>	<b>0.2 - 1.0 mL</b>	<b>2 - 10 mg</b>	<b>2 - 10 mL</b>
Adult Maximum Single Dose		10 mg	1.0 mL	10 mg	10 mL

Dosing Interval: **15 minutes** Adult **Max # of Doses: 4**

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

# Analgesia Medical Directive - Adult & Pediatric

## FentaNYL Dosing Guide

Route: Intravenous or Intranasal Supplied: 100 mcg in 2 mL *Intranasal Max Fluid : 1 mL per nare Use 1 mL Syringe, undiluted Maximum Pediatric Dosage: up to 1 mcg/kg (administer in divided doses)				
<b>⚠ Mandatory Provincial Patch Point for Children &lt; 12 years old</b>				
Age	Weight	Maximum Dose	Calculated Volume	Volume to administer (rounded)
Neonate	3 kg	3 mcg	0.06mL	0.05mL
<1	6 kg	6 mcg	0.12mL	0.1mL
1	12 kg	12 mcg	0.24 mL	0.2 mL
2	14 kg	14 mcg	0.28 mL	0.3 mL
3	16 kg	16 mcg	0.32 mL	0.3 mL
4	18 kg	18 mcg	0.36 mL	0.4 mL
5	20 kg	20 mcg	0.40 mL	0.4 mL
6	22 kg	22 mcg	0.44 mL	0.4 mL
7	24 kg	24 mcg	0.48 mL	0.5 mL
8	26 kg	26 mcg	0.52 mL	0.5 mL
9	28 kg	28 mcg	0.56 mL	0.6 mL
10	30 kg	30 mcg	0.60 mL	0.6 mL
11	32 kg	32 mcg	0.64 mL	0.6 mL
Youth* (12-17)	34 kg	34 mcg	0.68 mL	0.7 mL
	40 kg	40 mcg	0.80 mL	0.8 mL
	45 kg	45 mcg	0.90 mL	0.9 mL
	50 kg	50 mcg	1.0 mL	1.0 mL
	55 kg	55 mcg	1.1 mL*	1.1 mL*
	60 kg	60 mcg	1.2 mL*	1.2 mL*
	65 kg	65 mcg	1.3 mL*	1.3 mL*
	70 kg	70 mcg	1.4 mL*	1.4 mL*
	75 kg	75 mcg	1.5 mL*	1.5 mL*
Pediatric Maximum Single Dose*		75 mcg	1.5 mL*	1.5 mL*
Adults ≥ 18 years		25 – 75 mcg	0.50 -1.5 mL*	0.50 -1.5 mL*
Adult Maximum Single Dose		75 mcg	1.5 mL*	1.5 mL*

# Procedural Sedation Medical Directive – **AUXILIARY**

*An Advanced Care Paramedic may provide the treatment prescribed in this AUXILIARY Medical Directive if authorized.*

## INDICATIONS

Post-intubation; **OR**  
Transcutaneous pacing.

## CONDITIONS

### **fentaNYL**

AGE: ≥18 years  
LOA: N/A  
HR: N/A  
RR: ≥10/min\*  
SBP: Normotension  
Other: N/A

### **midazolam**

AGE: ≥18 years  
LOA: N/A  
HR: N/A  
RR: ≥10/min\*  
SBP: Normotension  
Other: N/A

**\*Non-intubated patients only**

## CONTRAINDICATIONS

### **fentaNYL**

Allergy or sensitivity to fentaNYL

### **midazolam**

Allergy or sensitivity to midazolam

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **fentaNYL**

	Route
	IV/IO/CVAD/IN
<i>Dose</i>	25-75 mcg
<i>Max. single dose</i>	75 mcg
<i>Dosing interval</i>	5 min
<i>Max. total dose</i>	150 mcg

Consider **midazolam**

	Route
	IV/IO/CVAD/IN
<i>Dose</i>	Up to 0.1 mg/kg
<i>Max. single dose</i>	5 mg
<i>Dosing interval</i>	5 min
<i>Max. total dose</i>	10 mg

## CLINICAL CONSIDERATIONS

Consider lower dose of medication in elderly and lighter weight individuals.

Consider quantitative EtCO<sub>2</sub> monitoring once the patient has been sedated.

## Combative Patient Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

### INDICATIONS

Combative **OR** violent or agitated behavior that requires sedation for patient safety.

### CONDITIONS

#### midazolam

AGE: ≥18 years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### ketamine

AGE: ≥18 years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected excited delirium /  
severe violent psychosis

### CONTRAINDICATIONS

#### midazolam

Allergy or sensitivity to midazolam

#### ketamine

Allergy or sensitivity to ketamine

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **midazolam**

Route	Age
	≥18 years
	IV/ IM / IN
<i>Dose</i>	Up to 0.1 mg/kg
<i>Max. single dose</i>	5 mg
<i>Dosing interval</i>	5 min
<i>Max. total dose</i>	10 mg
<i>Max. # doses</i>	N/A

Consider **ketamine**

Route	Age	Age
	≥18 years to <65 years	≥65 years
	IM	IM
<i>Dose</i>	5 mg/kg	3 mg/kg
<i>Max. single dose</i>	500 mg	300 mg
<i>Dosing interval</i>	N/A	N/A
<i>Max. # doses</i>	1	1



## CLINICAL CONSIDERATIONS

Reversible causes of combative, violent or agitated behaviours (e.g. hypoglycemia, hypoxia, hypovolemia) should be considered and treated (if possible) prior to treating with midazolam or ketamine.

Paramedics can administer a lower weight base dose (e.g. 0.05 mg/kg) of midazolam based on clinical judgment taking into consideration such as but not limited to, patient age, and degree of combativeness, and the level of suspicion of hypotension or hypoxia when unable to obtain vital signs.

Do not co-administer midazolam and ketamine unless direction received from BHP.

Consider quantitative EtCO<sub>2</sub> monitoring once the patient has been sedated.

If ketamine emergence reaction develops, a BHP patch is required if further sedation orders are required

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Nausea / Vomiting Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Nausea **OR** vomiting.

### CONDITIONS

#### ondansetron

AGE: N/A

Weight:  $\geq 25$  kg

LOA: Unaltered

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### dimenhyDRINATE

AGE: N/A

Weight:  $\geq 25$  kg

LOA: Unaltered

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### ondansetron

Allergy to ondansetron

Prolonged QT syndrome (known to patient)

Apomorphine use

#### dimenhyDRINATE

Allergy or sensitivity to dimenhyDRINATE or other antihistamines

Overdose on antihistamines or anticholinergics or tricyclic antidepressants

Co-administration of diphenhydramine

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **ondansetron**

	Weight ≥ 25 kg
	Route PO / IV* / IM*
Dose	4 mg
Max. single dose	4 mg
Dosing interval	N/A
Max. # of doses	1

\*IV / IM (if formulation is available and authorized)

Consider **dimenhyDRINATE**

	Weight ≥ 25 kg to < 50 kg	Weight ≥ 50 kg
	Route IV / IM	Route IV / IM
Dose	25 mg	** 25 or 50 mg
Max. single dose	25 mg	50 mg
Dosing interval	N/A	30 min
Max. # of doses	1	2
Max. Cumulative Dose	N/A	50 mg

\*\* If ondansetron is unavailable, assess the risks and benefits to patients ≥ 65 years old for dimenhyDRINATE administration. This may include an initial reduced dose of 25 mg

## CLINICAL CONSIDERATIONS

Prior to IV administration, dilute dimenhyDRINATE (concentration of 50 mg/1 ml) 1:9 with Normal Saline or D5W. If administered IM do not dilute  
If a patient has received an antiemetic and has no relief of their nausea & vomiting symptoms after 30 minutes, the alternative antiemetic may be considered.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank

# Procedural

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Home Dialysis Emergency Disconnect Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Patient receiving home dialysis (hemo or peritoneal) and connected to dialysis machine and requires transport to the closest appropriate receiving facility;

#### AND

Patient is unable to disconnect ;

#### AND

There is no family member of caregiver who is available and knowledgeable in dialysis disconnect .

### CONDITIONS

#### Home Dialysis Emergency Disconnect

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### Home Dialysis Emergency Disconnect

N/A

## TREATMENT

Consider **Home Dialysis Emergency Disconnect**

## CLINICAL CONSIDERATIONS

Generally, an emergency disconnect kit with materials and instructions can be found hanging from the dialysis machine or nearby on the wall.

Ensure both the patient side and machine side of the connection are clamped before disconnecting and attaching end caps.

## Emergency Dialysis Disconnect Prompt Card

### Hemodialysis

- ▶ Clamp patient side tubing clamps
- ▶ Clamp machine side clamps
- ▶ Attach sterile Luer Lock caps to the ends of the patient tubing
- ▶ Disregard any alarms that may sound from the machine
- ▶ Secure patient tubing and cover with abdo pad

### Continuous Ambulatory Peritoneal Dialysis (CAPD)

- ▶ Close the twist clamp
- ▶ Clamp both the fill and drain bag tubing with clamps supplied in disconnect kits
- ▶ Screw a sterile Luer Lock on the patient side tubing
  - Snap a sterile Luer Lock on the machine side tubing
- ▶ Secure patient tubing and cover with abdo pad

### Automatic Peritoneal Dialysis (APD)

- ▶ Push "Stop" button on ADP machine
- ▶ Close the twist clamp
- ▶ Disconnect the patient tubing from the machine tubing
- ▶ Screw a sterile mini cap on the patient tubing
- ▶ Snap a mini cap on the machine tubing
- ▶ Secure patient tubing and cover with abdo pad



# Emergency Childbirth Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

## INDICATIONS

Pregnant patient experiencing labour; **OR**

Post-partum patient immediately following delivery and/or placenta.

## CONDITIONS

### Delivery

AGE: Childbearing years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Second stage labour **AND/OR**  
Imminent birth **AND/OR**  
Shoulder Dystocia **AND/OR**  
Breech Delivery **AND/OR**  
Prolapsed Cord

### Umbilical Cord Management

AGE: Childbearing years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Cord complications **OR**  
if neonatal or maternal  
resuscitation is required **OR**  
Due to transport  
considerations

### External Uterine Massage

AGE: Childbearing years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Post-placental delivery

### oxytocin

AGE: Childbearing years

LOA: N/A

HR: N/A

RR: N/A

SBP: < 160 mmHg

Other: Postpartum delivery  
**AND/OR**  
Placental delivery

## CONTRAINDICATIONS

### Delivery

N/A

### Umbilical Cord Management

N/A

### External Uterine Massage

Placenta not delivered

### oxytocin

Allergy or sensitivity to oxytocin  
Undelivered fetus  
Suspected or known pre-eclampsia  
with current pregnancy  
Eclampsia (seizures) with current  
pregnancy  
≥ 4 hours post placenta delivery

## TREATMENT

### Consider **delivery**

Position the patient and deliver neonate

### Consider **shoulder dystocia delivery**

Perform ALARM twice on scene. If successful; deliver neonate. If unsuccessful; transport to closest appropriate facility

### Consider **breech delivery**

HANDS OFF the breech. Allow neonate to deliver to umbilicus; consider carefully releasing the legs &amp; arms as they are delivered; otherwise hands off.

Once hairline is visible AND/OR 3 mins has passed since umbilicus was visualized attempt the Mauriceau Smellie-Veit maneuver.

If successful; deliver neonate. If unsuccessful; transport to closest appropriate facility.

Consider **prolapsed cord delivery**

If a cord prolapse is present, the fetal part should be elevated to relieve pressure on the cord. Assist the patient into a knee-chest position or exaggerated Sims position, and insert gloved fingers/hand into the vagina to apply manual digital pressure to the presenting part which is maintained until transfer of care in hospital.

Airway /  
Breath.Cardiac /  
Circula.Consider **umbilical cord management**

If a nuchal cord is present and loose, slip cord over the neonate's head. Only if a nuchal cord is tight and cannot be slipped over the neonate's head, clamp and cut the cord, encourage rapid delivery.

Following delivery of the neonate, the cord should be clamped and cut immediately if neonatal or maternal resuscitation is required. Otherwise, after pulsations have ceased (approximately 2-3 minutes), clamp the cord in two places and cut the cord.

LOC

Pain/  
Sed./  
NauseaConsider **external uterine massage**

Post placental delivery

Proced.

Consider **oxytocin**

	Route
	IM
<i>Dose</i>	10 units
<i>Max. single dose</i>	10 units
<i>Dosing interval</i>	N/A
<i>Max. # doses</i>	1

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

## CLINICAL CONSIDERATIONS

If the patient presents with limb-presentation, do not attempt to push the limb back into the vagina; discourage the patient from pushing, cover the limb using a dry sheet to maintain warmth, and initiate transport as per the *Load and Go Patient Standard* of the BLS PCS.

If labour is failing to progress, discourage the patient from pushing or bearing down during contractions.

If delivery has not occurred at scene within approximately ten minutes of initial assessment, consider transport in conjunction with the following:

a. Patient assessment findings:

- i. Lack of progression of labour;
- ii. Multiple births expected;
- iii. Neonate presents face up;
- iv. Pre-eclampsia;
- v. Presence of vaginal hemorrhage;
- vi. Premature labour;
- vii. Primip;

b. Distance to the closest appropriate receiving facility.

When the placenta is delivered, inspect it for wholeness, place in a plastic bag from the OBS kit, label it with the maternal patient's name and time of delivery, and transport it with the maternal or neonatal patient. Delivery of the placenta should not delay transport considerations/initiation.

# Central Venous Access Device Access Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

## INDICATIONS

Actual or potential need for intravenous medication **OR** fluid therapy.

### AND

IV access is unobtainable;

### AND

Cardiac arrest or pre-arrest state.

## CONDITIONS

### CVAD Access

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Patient has a CVAD with an accessible external lumen

## CONTRAINDICATIONS

### CVAD Access

Inability to confirm patency of CVAD line

Inability to flush or aspirate

Injury or suspected fracture proximal to the access site

Swelling of the involved limb

Bleeding at the insertion site

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT

Consider **CVAD access**

## CLINICAL CONSIDERATIONS

N/A

# Lateral Patellar Dislocation Medical Directive

## - *AUXILIARY*

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Patient with suspected lateral patellar dislocation.

### CONDITIONS

#### Patellar Reduction

AGE: ≥ 10 years to ≤ 50 years  
LOA: Unaltered  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: N/A

### CONTRAINDICATIONS

#### Patellar Reduction

High velocity trauma  
Direct knee trauma

### TREATMENT

Consider **Patellar Reduction**

With the patient in a seated position, gently extend the knee while lifting up on the patella and placing medial pressure to the edge of the patella.

The maximum number of attempts for Patellar Reduction per patient is 2

### CLINICAL CONSIDERATIONS

N/A

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank



# Palliative Care / Research

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



## PALLIATIVE CARE MEDICAL DIRECTIVE

An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized to a patient with a life limiting illness **OR** a patient nearing end of life, requiring management for the following symptoms: Pain, Dyspnea, Nausea/Vomiting, Hallucinations/Delirium/Agitation, and/or Terminal Congested Breathing (noisy breathing or excessive secretions)

### Patch

If a paramedic determines that the patient would benefit from any other management that is not included in this medical directive, a patch to a Base Hospital Physician is **necessary**.

### Management of Patients with Palliative Care Needs

Patients with palliative care needs may require a different approach to assessment and treatment that reflects their unique goals of care. Therefore, for this defined patient population, paramedics should consider prioritizing patient comfort and are not required to follow the described regimen of strict vital signs, cardiac monitoring and transport as directed in the Basic Life Support Patient Care Standard (BLS PCS). If patient transport is initiated, paramedics should consider usual care (vitals and monitoring) per the ALS and BLS PCS in conjunction with the patients' goals of care; they may also consider symptom treatments below if indicated.

### Medical Directive

This Medical Directive is written in six sections including five symptom-based sections (Pain, Dyspnea, Hallucinations/Delirium/Agitation, Nausea/Vomiting and Terminal Congested Breathing) as well as a Treat and Refer directive. Any of these directives can apply, individually or in combination, to a patient with palliative care needs. The Treat and Refer part of this directive can be applied even if no symptoms listed in the directive are present or treatments have not been provided. All patients who remain at home must be referred to their primary care physician or a palliative care team to ensure follow up of their presenting complaint.

When in doubt, consult/patch to a Base Hospital Physician (BHP).

## Pocket Orders

Orders received from the base hospital physician that are for the **same patient** on the **same shift administered by the same paramedic**. They are designed for the paramedic to alleviate repeat calling when seeing the patient multiple times in a shift. If a different paramedic attends to the same patient, a new patch is required.

## Subcutaneous Infusion Device

A paramedic may insert a subcutaneous infusion device providing they have been trained and authorized **and** their employer has approved the use of this device and policies are in place supporting this technique

# Pain Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

## INDICATIONS

A patient with a life limiting illness **OR** a patient nearing end of life, requiring symptom management for pain.

## CLINICAL CONSIDERATIONS

- If the patient is opioid naïve, the lower range of doses should be used.
- If a patient is already on a regular opiate, the same opiate should be used. If the patient is on a regular opiate regimen that does not include either morphine or hydromorphone paramedics should confirm with a Base Hospital Physician prior to administering morphine or hydromorphone.
- If pain relief has not been achieved after maximum dosing, paramedics must consult with a Base Hospital Physician for further orders.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## CONDITIONS

### Morphine

AGE: ≥18  
LOA: N/A  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: N/A

### HYDROmorphine

AGE: ≥18  
LOA: N/A  
HR: N/A  
RR: N/A  
SBP: N/A  
Other: N/A

## CONTRAINDICATIONS

### Morphine

Allergy to morphine

### HYDROmorphine

Allergy to  
hydromorphine

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

### Consider Morphine

	Route
	SC/IV/CVAD
<i>Dose</i>	2-10mg
<i>Max. single dose</i>	10mg
<i>Dosing interval</i>	15 min
<i>Max. total dose</i>	2

### Consider HYDROmorphine

	Route
	SC/IV/CVAD
<i>Dose</i>	0.5 – 2mg
<i>Max. single dose</i>	2mg
<i>Dosing interval</i>	15 min
<i>Max. total dose</i>	2

## Dyspnea Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

A patient with a life limiting illness **OR** a patient nearing end of life, requiring symptom management for dyspnea.

### CLINICAL CONSIDERATIONS

- If patient is opioid naïve the lower range of doses should be considered.
- If the patient is already on a regular opiate, the same opiate should be used. If the patient is on a regular opiate regimen that does not include either morphine or hydromorphone, paramedics should confirm with a Base Hospital Physician prior to administering morphine or hydromorphone.
- If dyspnea has not been relieved after maximum dosing, paramedics must consult with a Base Hospital Physician for further orders.
- Salbutamol should only be used in patients whose dyspnea is **accompanied by wheezing or a history of response to bronchodilators.**

### CONDITIONS

Morphine	HYDROmorphine	Salbutamol
AGE: ≥18	AGE: ≥18	AGE: ≥18
LOA: N/A	LOA: N/A	LOA: N/A
HR: N/A	HR: N/A	HR: N/A
RR: N/A	RR: N/A	RR: N/A
SBP: N/A	SBP: N/A	SBP: N/A
Other: N/A	Other: N/A	Other: For dyspnea with suspected bronchoconstriction

## CONTRAINDICATIONS

**Morphine**

Allergy to morphine

**HYDROmorphine**Allergy to  
HYDROmorphine**Salbutamol**

Allergy to salbutamol

## TREATMENT

*Patient • Drug • Dose • Route • Time.*Consider **Morphine**

	Route
	SC/IV/CVAD
<i>Dose</i>	2-10mg
<i>Max. single dose</i>	10mg
<i>Dosing interval</i>	15 min
<i>Max. total dose</i>	2

Consider **HYDROmorphine**

	Route
	SC/IV/CVAD
<i>Dose</i>	0.5– 2mg
<i>Max. single dose</i>	2mg
<i>Dosing interval</i>	15 min
<i>Max. total dose</i>	2

Consider **Salbutamol**:

	Weight <25 kg		Weight ≥25 kg	
	Route MDI*	Route NEB	Route MDI*	Route NEB
<i>Dose</i>	Up to 600 mcg (6 puffs)	2.5 mg	Up to 800 mcg (8 puffs)	5 mg
<i>Max. single dose</i>	600 mcg	2.5 mg	800 mcg	5 mg
<i>Dosing interval</i>	5-15 min. PRN	5-15 min. PRN	5-15 min. PRN	5-15 min. PRN
<i>Max. # of doses</i>	3	3	3	3

\* 1 puff=100mcg

## Nausea/Vomiting Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

A patient with a life limiting illness **OR** a patient nearing end of life, requiring symptom management for nausea/vomiting.

### CLINICAL CONSIDERATIONS

- DimenhyDRINATE is rarely used in the palliative care population as it can cause delirium, increase drowsiness, and does not target the appropriate receptors to control the nausea in most patients. It should only be used in patients with contraindications to haloperidol where ondansetron cannot be used.

### CONDITIONS

Haloperidol	Ondansetron	DimenhyDRINATE
AGE: ≥18	AGE: ≥18	AGE: ≥18
LOA: N/A	LOA: N/A	LOA: N/A
HR: N/A	HR: N/A	HR: N/A
RR: N/A	RR: N/A	RR: N/A
SBP: N/A	SBP: N/A	SBP: N/A
Other: N/A	Other: contraindication to haloperidol	Other: contraindication to haloperidol AND ondansetron

## CONTRAINDICATIONS

**Haloperidol**

Allergy to haloperidol

Known Parkinson's  
or Lewy Body  
DementiaNeuroleptic  
Malignant Syndrome**Ondansetron**Allergy to  
ondansetron**DimenhyDRINATE**Allergy to  
dimenhyDRINATE or  
other antihistaminesOverdose on  
antihistamines or  
anticholinergics of  
tricyclic  
antidepressants

## TREATMENT

*Patient • Drug • Dose • Route • Time.*Consider **Haloperidol**

	Route
	SC/IV/CVAD
<i>Dose</i>	0.5-1mg
<i>Max. single dose</i>	1 mg
<i>Dosing interval</i>	30 min
<i>Max. total dose</i>	2

Consider **Ondansetron**

	Route
	PO/IV/SC/CVAD
<i>Dose</i>	4 mg
<i>Max. single dose</i>	4 mg
<i>Dosing interval</i>	N/A
<i>Max. total dose</i>	1

Consider **DimenhyDRINATE**

	Route
	SC/IV/CVAD
<i>Dose</i>	25-50mg
<i>Max. single dose</i>	50mg
<i>Dosing interval</i>	N/A
<i>Max. total dose</i>	1



## Hallucinations/Delirium/ Agitation Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

Airway /  
Breath.

### INDICATIONS

A patient with a life limiting illness **OR** a patient nearing end of life, requiring symptom management for hallucinations, delirium or agitation.

Cardiac /  
Circula.

### CLINICAL CONSIDERATIONS

- Haloperidol should be used as the first line agent for the treatment of hallucinations, delirium or agitation. Midazolam can be used in patients with contraindications to haloperidol.
- The patient receiving an approach to palliative care may show signs of **opioid toxicity**. Paramedics need to be cognizant of such conditions. In such cases patients may display signs of agitation, urticaria, myoclonus, hyperalgesia, pin point pupils. This can sometimes be interpreted as further pain. Consideration should be given to limit any further opioid dosing.
- If hallucinations, delirium or agitation has not been relieved after maximum dosing, paramedics must consult with a Base Hospital Physician for further orders.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

### CONDITIONS

Haloperidol	Midazolam
AGE: ≥18	AGE: ≥18
LOA: N/A	LOA: N/A
HR: N/A	HR: N/A
RR: N/A	RR: N/A
SBP: N/A	SBP: N/A
Other: N/A	Other: N/A

Chemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

## CONTRAINDICATIONS

### Haloperidol

Allergy to haloperidol

Known Parkinson's  
or Lewy Body  
DementiaNeuroleptic  
Malignant Syndrome

### Midazolam

Allergy to midazolam

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

### Consider **Haloperidol**

	Route
	SC/IV/CVAD
<i>Dose</i>	0.5-1mg
<i>Max. single dose</i>	1mg
<i>Dosing interval</i>	30 min
<i>Max. total dose</i>	2

### Consider **Midazolam**

	Route
	SC/IV/CVAD
<i>Dose</i>	0.5-2mg
<i>Max. single dose</i>	2mg
<i>Dosing interval</i>	30 min
<i>Max. total dose</i>	2

# Terminal Congested Breathing Medical Directive

An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.

## INDICATIONS

A patient with a life limiting illness **OR** a patient nearing end of life, requiring symptom management for terminal congested breathing (excessive secretions or noisy breathing).

Intended use of this medication is for a patient with a decreased level of consciousness in their last hours of life

## CLINICAL CONSIDERATIONS

- Patient repositioning and gentle turning of the head to the side can be done instead of medication however suction of the oropharynx **is not** appropriate as it will likely cause discomfort and a gag reflex.

## CONDITIONS

### Glycopyrrolate

AGE:  $\geq 18$

LOA: GCS 3-6

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### Atropine

AGE:  $\geq 18$

LOA: GCS 3-6

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## CONTRAINDICATIONS

### Glycopyrrolate

Allergy to  
glycopyrrolate

### Atropine

Allergy to atropine

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

### Consider **Glycopyrrolate**

	Route
	SC/IV/CVAD
<i>Dose</i>	0.4mg
<i>Max. single dose</i>	0.4mg
<i>Dosing interval</i>	N/A
<i>Max. total dose</i>	1

### Consider **Atropine**

	Route
	SC
<i>Dose</i>	0.4mg
<i>Max. single dose</i>	0.4mg
<i>Dosing interval</i>	N/A
<i>Max. total dose</i>	1

## Intravenous Cannulation Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

A patient with a life limiting illness **OR** a patient nearing end of life, requiring parenteral hydration or venous access.

### CLINICAL CONSIDERATIONS

- A follow up plan should be in place to ensure ongoing management of the intravenous line (such as follow up by MRP (most responsible practitioner) or community paramedic).
- A period of observation is recommended after the administration of any fluid if this patient is not transported to ensure adequate response and no unexpected immediate adverse effects.
- When a paramedic is requested to start an IV solely for the purpose of MAID (Medical Assistance in Dying) by the patient's care team, the paramedic has the right to refuse this request.

### CONDITIONS

#### IV Cannulation

AGE: ≥18

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## CONTRAINDICATIONS

### IV Cannulation

N/A

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider IV Cannulation

## Subcutaneous Line Placement Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

A patient with a life limiting illness **OR** a patient nearing end of life, requiring symptom management and parenteral administration of palliative care symptom relief medications is clinically indicated AND It is expected that more than one medication administration will be required

### CLINICAL CONSIDERATIONS

- A follow up plan should be in place to ensure ongoing management of the subcutaneous line (such as follow up by MRP or community paramedic).
- A period of observation is recommended after the administration of any medication if the patient is not transported to ensure adequate response and no unexpected immediate adverse effects.
- Adverse events after insertion include pain at the site (from the irritation of the drug or the injection was fast, this is prevented by injecting the drug slowly). If pain remain then the needle may be pulled back into the intradermal space (put a folded 2x2 gauze under the butterfly wings to elevate the needle to 45 degrees. If pain persist, then you need to change needle).

### CONDITIONS

#### Subcutaneous Line Placement

AGE:  $\geq 18$

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## CONTRAINDICATIONS

### Subcutaneous Line Placement

N/A

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider Subcutaneous Line Placement



## Treat and Refer Medical Directive

*An Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.*

### INDICATIONS

Symptoms improved to patient/Substitute Decision Maker (SDM) satisfaction

And

After informed discussion of goals of care, patient/SDM preference is to remain at home

And

After DNR and/or previous goals of care discussion with primary care provider

### CLINICAL CONSIDERATIONS

- A period of observation is recommended after the administration of any medication if the patient is not transported to ensure adequate response and no unexpected immediate adverse effects
- Transport should be considered if there is strong suspicion of reversible causes including but not limited to:
  - Complete bowel obstruction with no prior history of same
  - New Spinal Cord Compression
  - New Superior Vena Cava (SVC) Obstruction
  - Airway obstruction
  - Suspected new pathologic fracture

If the patient does not meet the treat and refer conditions, paramedics should consider consulting BHP, follow the patient refusal standard and document appropriately

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

---

**CONDITIONS****Treat and Refer**AGE:  $\geq 18$ 

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

---

**CONTRAINDICATIONS****Treat and Refer**

Concerns of patient abuse or neglect

Patient/SDM cannot demonstrate decision-making capacity based on the Aid to Capacity Evaluation Tool

Uncontrolled or new seizures

---

**TREATMENT*****Patient • Drug • Dose • Route • Time.***

Paramedics may assess and/or treat the patient according to this medical directive and, in collaboration with the patient/SDM, honour wishes to remain at home (treat and refer). Paramedics will notify the patients' palliative care team where the patient will remain at home to ensure follow up for their presenting complaint.

If the patient is not being followed by a palliative care team, the paramedic will ensure follow-up by a Community Agency or the Primary Care Physician.

Intentionally Left Blank

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

# The PRIME Trial Medical Directive

A Advanced Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.

## Indications

Pediatric non-traumatic cardiac arrest

## Conditions

CPR	
<b>Age</b>	≥ 24 hours to 17 years
<b>LOA</b>	Altered
<b>HR</b>	N/A
<b>RR</b>	N/A
<b>SBP</b>	N/A
<b>Other</b>	Performed in 2 minute intervals

Manual Defibrillation	
<b>Age</b>	≥ 24 hours to 17 years
<b>LOA</b>	Altered
<b>HR</b>	N/A
<b>RR</b>	N/A
<b>SBP</b>	N/A
<b>Other</b>	VF OR pulseless VT

AED Defibrillation	
<b>Age</b>	≥ 24 hours to 17 years
<b>LOA</b>	Altered
<b>HR</b>	N/A
<b>RR</b>	N/A
<b>SBP</b>	N/A
<b>Other</b>	Defibrillation indicated

Epinephrine Autoinjector	
<b>Age</b>	≥ 24 hours to 17 years
<b>LOA</b>	Altered
<b>HR</b>	N/A
<b>RR</b>	N/A
<b>SBP</b>	N/A
<b>Other</b>	N/A

## Contraindications

### CPR

Obviously dead as per BLS PCS

Meet conditions of Do Not  
Resuscitate (DNR) Standard

### Manual Defibrillation

Rhythms other than VF or pulseless  
VT

### AED Defibrillation

Non-shockable rhythm

### Epinephrine Autoinjector

Allergy or sensitivity to epinephrine

## Treatment

Consider CPR as described in the BLS PCS

Consider manual defibrillation (if available and authorized)

	Age	Age
	≥ 24 hours to < 8 years	≥ 8 years to 17 years
<b>Dose</b>	1 defibrillation	1 defibrillation
<b>Initial dose</b>	2 J/kg	As per BH / manufacturer
<b>Subsequent dose(s)</b>	4 J/kg	As per BH / manufacturer
<b>Dosing interval</b>	2 min	2 min
<b>Max. # of doses</b>	N/A	N/A

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

## Consider AED defibrillation (if not using manual defibrillation)

	Age		Age
	≥ 24 hours to < 8 years		≥ 8 years to 17 years
	With Pediatric Attenuator Cable	Without Pediatric Attenuator Cable	N/A
<b>Dose</b>	1 defibrillation	1 defibrillation	1 defibrillation
<b>Max. single dose</b>	As per BH / manufacturer	As per BH / manufacturer	As per BH / manufacturer
<b>Dosing interval</b>	2 min	2 min	2 min
<b>Max. # of doses</b>	N/A	N/A	N/A

## Consider EPINEPHrine Preload (if available)

	Route IM				
	Weight				
	≥ 3 kg to < 5 kg	≥ 5 kg to < 10 kg	≥ 10 kg to < 20 kg	≥ 20 kg to < 30 kg	≥ 30 kg
<b>Dose</b>	0.3 mg	0.5 mg	1 mg	2 mg	3 mg
<b>Total # of injections</b>	1	1	1	1	1
<b>Dosing interval</b>	N/A	N/A	N/A	N/A	N/A
<b>Max. # of doses</b>	1	1	1	1	1

Consider EPINEPHrine Autoinjector (if available)

Route				
IM				
	Weight			
	≥ 3 kg to < 5 kg	≥ 5 kg to < 10 kg	≥ 10 kg to < 20 kg	≥ 20 kg
<b>Dose</b>	0.3 mg	0.5 mg	1 mg	2 mg
<b>IM autoinjector used</b>	0.3 mg	0.5 mg	0.5 mg	0.5 mg
<b>Total # of injections</b>	1	1	2	4
<b>Dosing interval</b>	N/A	N/A	N/A	N/A
<b>Max. # of doses</b>	1	1	1	1

## Clinical Considerations

IM epinephrine should be administered as soon as possible once cardiac arrest identified and CPR initiated.

Continue standard care as per the Medical Cardiac Arrest Directive once IM epinephrine has been administered.

IV/IO epinephrine can be administered as soon as feasible after the initial IM epinephrine dose as per the Medical Cardiac Arrest Directive.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank





# Chemical Exposure

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES

## Cyanide Exposure Medical Directive – AUXILIARY CHEMICAL EXPOSURE

*An Advanced Care Paramedic may provide the treatment prescribed in this AUXILIARY Medical Directive if authorized.*

### INDICATIONS

Suspected exposure to cyanide

#### AND

Cardiac arrest; **OR**

Altered level of awareness; **OR**

Hypotension.

### CONDITIONS

#### Sodium thiosulfate 25%

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### hydroxocobalamin

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### Sodium thiosulfate 25%

Allergy or sensitivity to Sodium Thiosulfate 25%

#### hydroxocobalamin

Allergy or sensitivity to Hydroxocobalamin

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **sodium thiosulfate 25%**

Route	Age < 18 years	Age ≥ 18 years
	IV / IO / CVAD	IV / IO / CVAD
Dose	400 mg/kg or 1.6mL/kg over 15 min	12.5g (50 ml of 25% solution) over 15 min
Max. single dose	12.5g (50 ml of 25% solution)	12.5g (50 ml of 25% solution)
Dosing interval	N/A	N/A
Max. # doses	1	1



**Mandatory Provincial Patch Point**



Patch to BHP for authorization to proceed with the administration of hydroxocobalamin in cases of “suspected” cyanide toxicity.

Consider **hydroxocobalamin (if not using sodium thiosulfate 25%)**

Route	Age < 18 years	Age ≥ 18 years
	IV / IO / CVAD	IV / IO / CVAD
Dose	70 mg/kg over 30 min	5 g over 15 - 30 min
Max. single dose	5 g	5 g
Dosing interval	N/A	N/A
Max. # doses	1	1

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## CLINICAL CONSIDERATIONS

Hydroxocobalamin must be reconstituted with 200 ml normal saline prior to use.

### Hydroxocobalamin Dosing Chart

Weight (kg)	Dose	Concentration	Volume of Administration
<b>5</b>	350 mg	25 mg/ml	14 ml
<b>10</b>	700 mg	25 mg/ml	28 ml
<b>15</b>	1050 mg	25 mg/ml	42 ml
<b>20</b>	1400 mg	25 mg/ml	56 ml
<b>25</b>	1750 mg	25 mg/ml	70 ml
<b>30</b>	2100 mg	25 mg/ml	84 ml
<b>35</b>	2450 mg	25 mg/ml	98 ml
<b>40</b>	2800 mg	25 mg/ml	112 ml
<b>≥ 41</b>	5g	25 mg/ml	200 ml

## Hydrofluoric (HF) Acid Exposure Medical Directive - AUXILIARY CHEMICAL EXPOSURE

*An Advanced Care Paramedic may provide the treatment prescribed in this AUXILIARY Medical Directive if authorized.*

### INDICATIONS

Exposure to vapour and/or liquid hydrofluoric acid (HF);

#### AND

Exhibits signs and symptoms of HF acid toxicity.

### CONDITIONS

#### calcium gluconate

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

#### Topical Anaesthetic Eye Drops

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### calcium gluconate

Allergy or sensitivity to calcium gluconate

#### Topical Anaesthetic Eye Drops

Allergy or sensitivity to local anaesthetics

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT



*Patient • Drug • Dose • Route • Time.*

Consider **calcium gluconate**

	Inhalation exposure Concentration 10% solution	Skin exposure Concentration 2.5% gel
Route	NEB	TOP
Dose	100 mg	N/A
Max. single dose	100 mg	N/A
Dosing interval	N/A	Immediate
Max. # of doses	1	N/A

Consider **topical anaesthetic eye drops**

	Eye exposure
Route	TOP
Dose	2 gtts / eye
Max. single dose	2 gtts / eye
Dosing interval	10 min
Max. # of doses	N/A

## CLINICAL CONSIDERATIONS

For skin contact, ensure thorough irrigation prior to treatment.

For eye exposure remove patient's contact lenses, if applicable, prior to initiating treatment. Use anaesthetic eye drops for comfort and then irrigate eyes with normal saline for at least 15 minutes.

Administration of topical anesthetic eye drops should not delay the initiation of eye irrigation

Nebulizers typically require 2 to 3 mls to ensure appropriate medication administration.

## Adult Nerve Agent Exposure Medical Directive – AUXILIARY CHEMICAL EXPOSURE

*An Advanced Care Paramedic may provide the treatment prescribed in this AUXILIARY Medical Directive if authorized.*

### INDICATIONS

Exposure to a known or suspected nerve agent;

#### AND

Signs and symptoms of a cholinergic crisis

### CONDITIONS

#### atropine

AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected cholinergic crisis

#### Moderate Exposure

Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure

#### Severe Exposure

Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

**pralidoxime**AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected cholinergic crisis

**Moderate Exposure**

Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure

**Severe Exposure**

Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea

**diazepam**AGE:  $\geq 18$  years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected cholinergic crisis

**Moderate Exposure**

Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure

**Severe Exposure**

Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea



midazolam	
AGE:	≥ 18 years
LOA:	N/A
HR:	N/A
RR:	N/A
SBP:	N/A
Other: Suspected cholinergic crisis	
<b>Moderate Exposure</b>	
Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure	
<b>Severe Exposure</b>	
Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea	

## CONTRAINDICATIONS

<b>atropine</b> Allergy or sensitivity to atropine	<b>pralidoxime</b> Allergy or sensitivity to pralidoxime
<b>diazepam</b> Allergy or sensitivity to diazepam	<b>midazolam</b> Allergy or sensitivity to midazolam Use / Availability of diazepam

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **atropine**

Route	Moderate exposure IM	Severe Exposure IM
Initial Dose	2 mg	6 mg
Additional doses	2 mg	6 mg
Dosing interval	5 min.	5 min.
Max. # of doses	N/A	N/A

Consider **pralidoxime**

Route	Moderate exposure IM	Severe Exposure IM
Initial Dose	600 mg	1,800 mg
Additional doses	600 mg	1,800 mg
Dosing interval	15 min.	60 min.
Max. # of doses	3	2

Consider **diazepam**

Route	Moderate exposure IM
<i>Dose</i>	10 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

Consider **midazolam** (if not using diazepam)

Route	Moderate exposure IM
<i>Initial Dose</i>	10 mg
<i>Dosing interval</i>	5 min.
<i>Max. # of doses</i>	2

## CLINICAL CONSIDERATIONS

Atropine should be administered prior to airway interventions if secretions are copious.

Pralidoxime should be given as soon as possible after the administration of atropine.

Subsequent doses of atropine are intended for patients showing signs of bronchial secretions and may be repeated as indicated until airway secretions are controlled.

Decontamination procedures must be integrated with antidote administration.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Pediatric Nerve Agent Exposure Medical Directive – AUXILIARY CHEMICAL EXPOSURE

*An Advanced Care Paramedic may provide the treatment prescribed in this AUXILIARY Medical Directive if authorized.*

### INDICATIONS

Exposure to a known or suspected nerve agent;

#### AND

Signs and symptoms of a cholinergic crisis

### CONDITIONS

#### atropine

AGE: < 18 years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected cholinergic crisis

#### Moderate Exposure

Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure

#### Severe Exposure

Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea

**pralidoxime**

AGE: &lt; 18 years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected cholinergic crisis

**Moderate Exposure**

Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure

**Severe Exposure**

Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.**diazepam**

AGE: &lt; 18 years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected cholinergic crisis

**Moderate Exposure**

Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure

**Severe Exposure**

Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## midazolam

AGE: < 18 years

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: Suspected cholinergic crisis

### Moderate Exposure

Any one of the following: vomiting, diarrhea, bronchospasm or bronchial secretions, shortness of breath or any known liquid exposure

### Severe Exposure

Signs and symptoms of a moderate exposure and any one of the following: decreased LOA, paralysis, seizure or apnea

## CONTRAINDICATIONS

### atropine

Allergy or sensitivity to atropine

### pralidoxime

Allergy or sensitivity to pralidoxime

### diazePAM

Allergy or sensitivity to diazePAM

### midazolam

Allergy or sensitivity to midazolam

Use / Availability of diazePAM

## TREATMENT



**Patient • Drug • Dose • Route • Time.**

Consider **atropine**

	Moderate and Severe exposure	Moderate and Severe Exposure	Moderate Exposure	Severe Exposure
	Weight < 10 kg	Weight ≥ 10 kg to < 40 kg	Weight ≥ 40 kg	Weight ≥ 40 kg
Route	IM	IM	IM	IM
Dose	0.5 mg	1 mg	2 mg	6 mg
Max. single dose	0.5 mg	1 mg	2 mg	6 mg
Dosing interval	5 min.	5 min.	5 min.	5 min.
Max. # of doses	N/A	N/A	N/A	N/A

Consider **pralidoxime**

	Moderate exposure	Severe Exposure	Moderate Exposure	Severe Exposure
	Weight < 40 kg	Weight < 40 kg	Weight ≥ 40 kg	Weight ≥ 40 kg
Route	IM	IM	IM	IM
Dose	15 mg/kg	45 mg/kg	600 mg	1,800 mg
Max. single dose	600 mg	600 mg	600 mg	1,800 mg
Dosing interval	15 min.	60 min.	15 min.	60 min.
Max. # of doses	3	2	3	2

Consider **diazepam**

	Moderate exposure	Severe Exposure
	Weight	Weight
	< 50 kg	≥ 50 kg
Route	IM	IM
Dose	0.2 mg/kg	10 mg
Max. single dose	10 mg	10 mg
Dosing interval	N/A	N/A
Max. # of doses	1	1

Consider **midazolam** (if not using diazepam)

	Moderate exposure	Severe Exposure
	Weight	Weight
	< 50 kg	≥ 50 kg
Route	IM	IM
Dose	0.2 mg/kg	10
Max. single dose	10 mg	10 mg
Dosing interval	5 min.	5 min.
Max. # of doses	2	2

## CLINICAL CONSIDERATIONS

Consider using autoinjector for patients who are <50 kg with severe symptoms if there is any perceived delay to treatment.

Atropine should be administered prior to airway interventions if secretions are copious.

Pralidoxime should be given as soon as possible after the administration of atropine.

Subsequent doses of atropine are intended for patients showing signs of bronchial secretions and may be repeated as indicated until airway secretions are controlled.

Decontamination procedures must be integrated with antidote administration.



## Symptomatic Riot Agent Medical Directive – AUXILIARY CHEMICAL EXPOSURE

*An Advanced Care Paramedic may provide the treatment prescribed in this AUXILIARY Medical Directive if authorized.*

### INDICATIONS

Known or suspected exposure to a riot agent with signs and symptoms of a riot agent exposure.

### CONDITIONS

#### Topical anaesthetic eye drops

AGE: N/A

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

### CONTRAINDICATIONS

#### acetaminophen

Allergy or sensitivity to local anaesthetics

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## TREATMENT

Consider **topical anaesthetic eye drops**

Route	TOP
<i>Dose</i>	2 gtts / eye
<i>Max. single dose</i>	2 gtts / eye
<i>Dosing interval</i>	10 min
<i>Max. # doses</i>	N/A

## CLINICAL CONSIDERATIONS

For skin or mucous membrane contact, ensure thorough irrigation.  
For eye exposure, remove patient's contact lenses if applicable prior to initiating treatment. Use anaesthetic eye drops for comfort and then irrigate eyes with normal saline for at least 15 minutes.



## Medical References


ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



## ETCO<sub>2</sub> Waveforms

### Sudden loss waveform

- ET tube disconnected, dislodged, kinked or obstructed
- Loss of circulatory function



### Decreasing EtCO<sub>2</sub>

- ET tube cuff leak
- ET tube in hypopharynx
- Partial obstruction



### CPR Assessment

- Attempt to maintain minimum of 10 mmHg




### Sudden increase in EtCO<sub>2</sub>

- Return of spontaneous circulation (ROSC)



### Bronchospasm ("Shark-fin" appearance)


- Asthma
- COPD



### Hypoventilation



### Hyperventilation




### Decreased EtCO<sub>2</sub>

- Apnea
- Sedation



## 12 Lead ECG Placement



### PRECORDIAL LEADS:

- V1** - 4<sup>th</sup> intercostal space to the right of the sternum
- V2** - 4<sup>th</sup> intercostal space to the left of the sternum
- V3** - directly between leads V2 and V4
- V4** - 5<sup>th</sup> intercostal space at left midclavicular line
- V5** - level with lead V4 at left anterior axillary line
- V6** - level with lead V5 at left midaxillary line

### LIMB LEADS

- RA** - right forearm or wrist
- LA** - left forearm or wrist
- LL** - left lower leg
- RL** - right lower leg


### NOTE:

Refer to the Medical Directives for the clinical situations where a 12-Lead ECG should be considered. This may include patients experiencing cardiac ischemia, acute cardiogenic pulmonary edema, tachycardias, bradycardias, shortness of breath or upon ROSC.

## STEMI Anatomical Location

<b>I</b> Lateral	<b>aVR</b>	<b>V1</b> Septal	<b>V4</b> Anterior
<b>II</b> Inferior	<b>aVL</b> Lateral	<b>V2</b> Septal	<b>V5</b> Lateral
<b>III</b> Inferior	<b>aVF</b> Inferior	<b>V3</b> Anterior	<b>V6</b> Lateral

## 15-Lead ECG Placement



- V4 becomes **V4R** - fifth intercostal space at **right** midclavicular line (similar position as V4 but on right side of chest)
- V5 becomes **V8** - level with V6 at left midscapular line
- V6 becomes **V9** - level with V6 at left paravertebral line

### NOTE:

- Limb leads should be placed on the limbs and not on the chest
- Consider assessing V4R when the 12 Lead identifies an inferior STEMI or ST depression in any of the septal leads (V1/V2).
- Consider assessing V8 and V9 when the 12 lead shows ST depression in the precordial leads or the 12 lead appears 'normal'.
- ST elevation of  $\geq 1$  mm in V4R and inferior ST-elevation, suggests a Right Ventricular involvement.
- ST elevation of  $\geq 1$  mm or greater in V8 and V9 suggests Posterior MI.

## CPR Guidelines

	Recommendations		
Component	★ Adults	★ Children	★ Infants
Recognition	★★★ Check for responsiveness (for all ages) ★★★ No breathing or only gasping (ie: abnormal) ★★★ No pulse palpated within 10 seconds for all ages ★★ HR < 60 and signs of hypoperfusion		
CPR sequence	★★★ C-A-B		
Compression rate	★★★ 100-120/min		
Compression depth	★ 5.0 – 6.0 cm (2.0 - 2.4 inches)	★ At least 1/3 AP diameter ★ About 5 cm (2 inches)	★ At least 1/3 AP diameter ★ About 4 cm (1½ inches)
Chest wall recoil	★★★ Allow complete recoil between compressions Rotate compressors every 2 minutes		
Compression interruptions	★★★ Minimize interruptions in chest compressions Attempt to limit interruptions to < 10 seconds		
Airway	★★★ Head tilt-chin lift or where trauma is suspected, jaw thrust		
Compression-to-ventilation ratio (until advanced airway placed)	★ 30:2 1 or 2 rescuers	★★ 30:2 Single rescuer ★★ 15:2 2 HCP rescuers  Neonates: 3:1	
Ventilations with advnaced airway (HCP)	★★★ 1 breath every 6-8 seconds (8-10 breaths/min) Asynchronous with chest compressions About 1 second per breath without too much force Visible chest rise		
Defibrillation	★★★ Attach and use AED as soon as available. Minimize interuptions in CPR pre & post rhythm interpretation/defibrillation to < 10 seconds		

### CPR NOTES:

- ▶ Rate: 100-120 compressions/minute and allow full chest recoil.
- ▶ Switch person doing compressions every 2 minutes and focus on high quality CPR.
- ▶ Minimize interruptions to chest compressions at all times.
- ▶ Give ventilations over 1 second just to point of seeing chest rise.

**ADULTS:**

Non-intubated: ratio 30:2 as above.

Intubated: 10 ventilations per minute without interrupting chest compressions.

SGA inserted: 10 ventilations per minute without interrupting chest compressions

**PEDIATRICS (30 DAYS TO AGE 12):**

Non-intubated: ratio 15:2 as above.

Intubated: 8-10 ventilations per minute without interrupting chest compressions.

Ventilations for resp. arrest only, non-intubated: 12-20/min.

**NEONATE:**

Both non-intubated **AND** intubated 3:1 ratio as above.

**ETCO<sub>2</sub> IN CARDIAC ARREST**

- ▶ When a SGA or ETT is in place, the following concepts apply:
- ▶ Continuous waveform capnography is recommended in addition to clinical assessment as the most reliable method of confirming and monitoring correct placement of an endotracheal tube
- ▶ Waveform capnography should be used to confirm and monitor endotracheal tube and SGA placement at all times
- ▶ Studies on waveform capnography have shown nearly 100% sensitivity and 100% specificity in identifying correct endotracheal tube and SGA placement
- ▶ Using quantitative waveform capnography is recommended in patients to monitor CPR quality, optimize chest compressions, and detect ROSC during chest compressions or when rhythm check reveals an organized rhythm (in addition to pulse checks)
- ▶ If waveform capnography abruptly increases to a normal value (35 to 40 mm Hg or higher) and is sustained, this may represent ROSC; wait for the next rhythm check to check for a pulse (or stop sooner if the patient exhibits signs of life)
- ▶ An ETCO<sub>2</sub> < 10 mmHg in VSA patients after 20 minutes of ACLS have a very poor prognosis; and can be used with clinical factors for the BHP to determine if TOR is appropriate.




# Rule of Nines, Burn Percentage Chart

## PEDIATRIC



## ADULT



Advanced Trauma Life Support, 9<sup>th</sup> Edition 2012 ; The American College of Surgeons.

Medical References Rule of Nines, Burns Percentage Chart

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Intramuscular Injection

- ▶ An intramuscular (IM) injection is a parenteral medication administration route commonly used by paramedics. It involves injecting a pharmacological agent directly into muscle tissue. Muscle tissue is very vascular, and as a result IM injections tend to have a faster onset of action than subcutaneous administrations.
- ▶ Identify patient that meets criteria for an intramuscular medication administration (refer to the Medical Directives or BHP order).
- ▶ Ensure all the “rights” of medication administration have been met
- ▶ Confirm medication and dose with paramedic partner if available.
- ▶ Follow safe process for responsible medication administration.
- ▶ Landmark the intended injection site. Generally the deltoid and the vastus lateralis are easily accessible and appropriate sites for IM injections; however other sites may be appropriate and can be landmarked as per the diagram on the following page.
- ▶ Select the appropriate size and gauge needle.
- ▶ Cleanse the needle insertion site using aseptic technique.
- ▶ Prepare the appropriate medication and dose into the syringe and needle ensuring all air bubbles are removed prior to injection.
- ▶ Stretch the skin taut and use the “Z-track” technique to displace the skin and soft tissue. Insert the needle with syringe/medication at a 90 degree angle using a “dart style” motion. The Z-track method reduces the chance the medication will leak from the muscle into the subcutaneous tissue.
- ▶ Inject the correct dose of medication.
- ▶ Remove the needle and immediately dispose of it in the biohazard container.
- ▶ Apply gentle pressure to the site with a dry gauze. Do not rub or massage. Apply a band-aid if needed.

## Intramuscular Injection Sites




Figure 1 - Deltoid




Figure 2 – Vastus Lateralis




Figure 3 – Ventrogluteal

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Formulas

**NOTE:** The formulas below are for reference purposes only. Paramedics must refer to the Medical Directives and/or Base Hospital Physician patch orders for appropriate treatment options.

### IV FLOW RATE CALCULATION:

$$\text{gtt/min} = \frac{\text{Amount (mL) to be infused} \times \text{Drops per mL (gtt/mL) of administration set}}{\text{Total time of infusion (min)}}$$

### MEDICATION INFUSION RATE:

$$\text{mL/hr} = \frac{\text{Desired dose (mg/min)} \times 60 \text{ min/hr}}{\text{Drug concentration (mg/mL)}}$$

**Note:** Units must be consistent throughout the calculation. For example, the desired dose can be in mcg/min, as long as the concentration is also converted into mcg/mL.

### PEDIATRIC BODY WEIGHT:

**For use with children aged 1 to 10 years.**

(Age in years  $\times$  2) + 10 = Approximate child body weight in kg.

### OXYGEN TANK DURATION:


$$\text{Duration of flow (minutes)} = \frac{\text{Gauge pressure} - \text{Safe residual pressure}}{\text{Flow rate (L/min)}} \times \text{Cylinder factor}$$

Cylinder Factor: D-tank = 0.16; M-tank = 1.56

# Spinal Motion Restriction Standard

## Prompt Card

This prompt card provides a quick reference of the *Spinal Motion Restriction (SMR) Standard* contained in the *Basic Life Support Patient Care Standards* (BLS PCS). Please refer to the BLS PCS for the full standard.



## “Single Strength” DOPamine Dosing Guide

DOPamine INFUSION RATE (mL/hr or drops/min with a microdrip set)  
[Using an 800 mcg/ml ('single strength') solution]

Weig ht (kg)	Drip Rate (drops/min)				
	2 (mcg/kg/minu te)	5 (mcg/kg/minu te)	10 (mcg/kg/minu te)	15 (mcg/kg/minu te)	20 (mcg/kg/minu te)
5	1	2	4	6	8
10	2	4	8	11	15
15	2	6	11	17	23
20	3	8	15	23	30
25	4	9	19	28	38
30	5	11	23	34	45
35	5	13	26	39	53
40	6	15	30	45	60
45	7	17	34	51	68
50	8	19	38	56	75
55	8	21	41	62	83
60	9	23	45	68	90
65	10	24	49	73	98
70	11	26	53	79	105
75	11	28	56	84	113
80	12	30	60	90	120
85	13	32	64	96	128
90	14	34	68	101	135
95	14	36	71	107	143
100	15	38	75	113	150
105	16	39	79	118	158
110	17	41	83	124	165
115	17	43	86	129	173
120	18	45	90	135	180

## EPINEPHrine 1 mg/mL = 1:1000 IM

### Dosing Guide

*Dose (0.01 mg/kg) is rounded to the nearest 0.05mg  
Use a 1 mL syringe*

AGE	WEIGHT	DOSE (mg)	VOLUME (mL) (rounded)
3 months	5 kg	0.05 mg	0.05 mL
6 months	8 kg	0.08 mg	0.10 mL
9 months	10 kg	0.10 mg	0.10 mL
1 year	12 kg	0.12 mg	0.10 mL
2 years	14 kg	0.14 mg	0.15 mL
3 years	16 kg	0.16 mg	0.15 mL
4 years	18 kg	0.18 mg	0.20 mL
5 years	20 kg	0.20 mg	0.20 mL
6 years	22 kg	0.22 mg	0.20 mL
7 years	24 kg	0.24 mg	0.25 mL
8 years	26 kg	0.26 mg	0.25 mL
9 years	28 kg	0.28 mg	0.30 mL
10 years	30 kg	0.30 mg	0.30 mL
11 years	32 kg	0.32 mg	0.30 mL
12 years	34 kg	0.34 mg	0.35 mL
13 years	36 kg	0.36 mg	0.35 mL
14 years	38 kg	0.38 mg	0.40 mL
Adult	50 kg	0.50 mg	0.50 mL

## Analgesia Medical Directive - Adult & Pediatric Morphine Dosing Guide

Age	Weight	Route: Subcutaneous Pediatric dosage 0.05 mg/kg Supplied: 10 mg/mL Use 1 mL Syringe Undiluted			Route: Intravenous Pediatric dosage 0.05 mg/kg Supplied: 10 mg/mL Use 1 mL Syringe Diluted to 1 mg/mL		
		Dose	Calculated Volume	Volume To Administer (rounded)	Dose	Calculated Volume	Volume To Administer (rounded)
		<div><div></div><div>Mandatory Provincial Patch Point</div></div>					
		For patients < 12 years					
Neonate	3 kg	0.15 mg	0.015 mL	0.02 mL	0.15 mg	0.015 mL	0.02 mL
<1	6 kg	0.3 mg	0.03 mL	0.05 mL	0.3 mg	0.03 mL	0.3 mL
1	12 kg	0.6 mg	0.06 mL	0.05 mL	0.6 mg	0.6 mL	0.6 mL
2	14 kg	0.7 mg	0.07 mL	0.05 mL	0.7 mg	0.7 mL	0.7 mL
3	16 kg	0.8 mg	0.08 mL	0.10 mL	0.8 mg	0.8 mL	0.8 mL
4	18 kg	0.9 mg	0.09 mL	0.10 mL	0.9 mg	0.9 mL	0.9 mL
5	20 kg	1.0 mg	0.10 mL	0.10 mL	1.0 mg	1.0 mL	1.0 mL
6	22 kg	1.1 mg	0.11 mL	0.10 mL	1.1 mg	1.1 mL	1.0 mL
7	24 kg	1.2 mg	0.12 mL	0.1 mL	1.2 mg	1.2 mL	1.2 mL
8	26 kg	1.3 mg	0.13 mL	0.1 mL	1.3 mg	1.3 mL	1.4 mL
9	28 kg	1.4 mg	0.14 mL	0.1 mL	1.4 mg	1.4 mL	1.4 mL
10	30 kg	1.5 mg	0.15 mL	0.2 mL	1.5 mg	1.5 mL	1.6 mL
11	32 kg	1.6 mg	0.16 mL	0.2 mL	1.6 mg	1.6 mL	1.6 mL
		Supplied: 10 mg/mL Use 1 mL Syringe Undiluted			Supplied: 10 mg/mL Use 10 mL Syringe Diluted to 1 mg/mL		
Youth (12-17)	34 kg	1.7 mg	0.17 mL	0.2 mL	1.7 mg	1.7 mL	1.8 mL
	40 kg	2.0 mg	0.20 mL	0.2 mL	2.0 mg	2.0 mL	2.0 mL
	45 kg	2.25 mg	0.225 mL	0.2 mL	2.25 mg	2.25 mL	2.2 mL
	50 kg	2.5 mg	0.25 mL	0.3 mL	2.5 mg	2.5 mL	2.6 mL
	55 kg	2.75 mg	0.275 mL	0.3 mL	2.75 mg	2.75 mL	2.8 mL
	60 kg	3.0 mg	0.30 mL	0.3 mL	3.0 mg	3.0 mL	3.0 mL
	65 kg	3.25 mg	0.325 mL	0.3 mL	3.25 mg	3.25 mL	3.2 mL
	70 kg	3.5 mg	0.35 mL	0.4 mL	3.5 mg	3.5 mL	3.6 mL
	75 kg	3.75 mg	0.375 mL	0.4 mL	3.75 mg	3.75 mL	3.8 mL
	80 kg	4.0 mg	0.40 mL	0.4 mL	4.0 mg	4.0 mL	4.0 mL
	85 Kg	4.25 mg	0.425 mL	0.4 mL	4.25 mg	4.25 mL	4.2 mL
	90 kg	4.5 mg	0.45 mL	0.5 mL	4.5 mg	4.5 mL	4.6 mL
95 kg	4.75 mg	0.475 mL	0.5 mL	4.75 mg	4.75 mL	4.8 mL	
100 kg	5 mg	0.5 mL	0.5 mL	5.0 mg	5.0 mL	5.0 mL	
Pediatric Maximum Single Dose		5 mg	0.50 mL	0.5 mL	5.0 mg	5 mL	5 mL

Dosing Interval: 15 minutes Pediatric Max # of Doses: 4



## Analgesia Medical Directive - Adult & Pediatric Morphine Dosing Guide

		Supplied: 10 mg/mL Use 1 mL Syringe Undiluted		Supplied: 10 mg/mL Use 10 mL Syringe Diluted to 1 mg/mL	
<b>Adult</b>	<b>N/A</b>	2 - 10mg	0.2 - 1.0 mL	2 - 10 mg	2 - 10 mL
<b>Adult Maximum Single Dose</b>		10 mg	1.0 mL	10 mg	10 mL

Dosing Interval: **15 minutes**    Adult **Max # of Doses: 4**

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

## Analgesia Medical Directive - Adult & Pediatric

### FentaNYL Dosing Guide

Route: Intravenous or Intranasal

Supplied: 100 mcg in 2 mL

\*Intranasal Max Fluid : 1 mL per nare

Use 1 mL Syringe, undiluted

Maximum Pediatric Dosage: up to 1 mcg/kg (administer in divided doses)



**Mandatory Provincial Patch Point for Children < 12 years old**

Age	Weight	Maximum Dose	Calculated Volume	Volume to administer (rounded)
Neonate	3 kg	3 mcg	0.06mL	0.05mL
<1	6 kg	6 mcg	0.12mL	0.1mL
1	12 kg	12 mcg	0.24 mL	0.2 mL
2	14 kg	14 mcg	0.28 mL	0.3 mL
3	16 kg	16 mcg	0.32 mL	0.3 mL
4	18 kg	18 mcg	0.36 mL	0.4 mL
5	20 kg	20 mcg	0.40 mL	0.4 mL
6	22 kg	22 mcg	0.44 mL	0.4 mL
7	24 kg	24 mcg	0.48 mL	0.5 mL
8	26 kg	26 mcg	0.52 mL	0.5 mL
9	28 kg	28 mcg	0.56 mL	0.6 mL
10	30 kg	30 mcg	0.60 mL	0.6 mL
11	32 kg	32 mcg	0.64 mL	0.6 mL
Youth* (12-17)	34 kg	34 mcg	0.68 mL	0.7 mL
	40 kg	40 mcg	0.80 mL	0.8 mL
	45 kg	45 mcg	0.90 mL	0.9 mL
	50 kg	50 mcg	1.0 mL	1.0 mL
	55 kg	55 mcg	1.1 mL*	1.1 mL*
	60 kg	60 mcg	1.2 mL*	1.2 mL*
	65 kg	65 mcg	1.3 mL*	1.3 mL*
	70 kg	70 mcg	1.4 mL*	1.4 mL*
	75 kg	75 mcg	1.5 mL*	1.5 mL*
Pediatric Maximum Single Dose*		75 mcg	1.5 mL*	1.5 mL*
Adults ≥ 18 years		25 – 75 mcg	0.50 -1.5 mL*	0.50 -1.5 mL*
Adult Maximum Single Dose		75 mcg	1.5 mL*	1.5 mL*

\*for pediatric dosing, consider administering in divided doses of one-third to one-half and titrate to effect similar to adult dosing.

# Seizure Medical Directive Dosing Guide

## Midazolam Dosing Guide

Age	Weight	Route: IM/IN/Buccal			Route: IV/IO		
		Dose: 0.2 mg/kg Supplied: 10 mg/2 mL Use 1 mL syringe Undiluted			Dose: 0.1 mg/kg Supplied: 10 mg/2 mL Use 10 mL syringe diluted to 1 mg/mL		
		Dose	Calculated Volume	Volume to Administer (rounded)	Dose	Actual Volume	Volume to Administer (rounded)
Neonate	3 kg	0.6 mg	0.12 mL	0.10 mL	0.3 mg	0.3 mL	0.4 mL
< 1	6 kg	1.2 mg	0.24 mL	0.25 mL	0.6 mg	0.6 mL	0.6 mL
1	12 kg	2.4 mg	0.48 mL	0.50 mL	1.2 mg	1.2 mL	1.2 mL
2	14 kg	2.8 mg	0.56 mL	0.55 mL	1.4 mg	1.4 mL	1.4 mL
3	16 kg	3.2 mg	0.64 mL	0.65 mL	1.6 mg	1.6 mL	1.6 mL
4	18 kg	3.6 mg	0.72 mL	0.70 mL	1.8 mg	1.8 mL	1.8 mL
5	20 kg	4.0 mg	0.80 mL	0.80 mL	2.0 mg	2.0 mL	2.0 mL
6	22 kg	4.4 mg	0.88 mL	0.90 mL	2.2 mg	2.2 mL	2.2 mL
		Supplied: 10 mg/2 mL Use 3 mL or 10 mL syringe Undiluted			Supplied: 10 mg/2 mL Use 10 mL syringe Diluted to 1 mg/mL		
7	24 kg	4.8 mg	0.96 mL	1.0 mL	2.4 mg	2.4 mL	2.4 mL
8	26 kg	5.2 mg	1.04 mL	1.0 mL	2.6 mg	2.6 mL	2.6 mL
9	28 kg	5.6 mg	1.12 mL	1.2 mL	2.8 mg	2.8 mL	2.8 mL
10	30 kg	6 mg	1.20 mL	1.2 mL	3.0 mg	3.0 mL	3.0 mL
11	32 kg	6.4 mg	1.28 mL	1.2 mL	3.2 mg	3.2 mL	3.2 mL
12	34 kg	6.8 mg	1.36 mL	1.4 mL	3.4 mg	3.4 mL	3.4 mL
	40 kg	8 mg	1.60 mL	1.6 mL	4.0 mg	4.0 mL	4.0 mL
	45 kg	9 mg	1.80 mL	1.8 mL	4.5 mg	4.5 mL	4.5 mL
Max	>50 kg	10 mg	2.00 mL	2.0 mL	5.0 mg	5.0 mL	5.0 mL

Note: Dosage administered can be calculated by the weight based calculation in the Medical Directive and/or by using the above chart. Administered dosage in the chart may be rounded to the nearest volume increment that can be accurately measured.

### Note:

Dosing for Adult Procedural Sedation: up to 0.1mg/kg (IV/IM/IN); max single dose 5mg; max 2 doses

Dosing for Adult Combative Patient up to 0.1mg/kg (IV/IO/CVAD/IN); max single dose 5mg; max total dose 10mg

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank



# Medication Information

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES

## Medication Information

### ACETAMINOPHEN

<b>CLASS</b>	Antipyretic and analgesic. Mild anti-inflammatory effects.
<b>ACTION</b>	Exact mechanism is not known. Rapidly absorbed through GI tract. Believed to raise the pain threshold.
<b>ONSET</b>	15 minutes and lasts up to 3 hours.
<b>METABOLISM</b>	At normal therapeutic dosages, primary hepatic metabolism. A toxic dose (as little as 4g daily) can cause hepatic cell necrosis. Oral administration is subject to first pass metabolism.

### ADENOSINE

<b>CLASS</b>	Antiarrhythmic
<b>ACTION</b>	Slows conduction time through the AV node, interrupting the re-entry pathways through the AV node, restoring normal sinus rhythm. Adenosine also causes coronary vasodilation and increases blood flow in normal coronary arteries with little to no increase in stenotic coronary arteries; thallium-201 uptake into the stenotic coronary arteries will be less than that of normal coronary arteries revealing areas of insufficient blood flow.
<b>ONSET</b>	Rapid
<b>HALF-LIFE</b>	< 10 seconds
<b>METABOLISM</b>	Blood and tissue.

### AMIODARONE

<b>CLASS:</b>	Antiarrhythmic (Class I, II, III, and IV)
<b>ACTION:</b>	Blocks sodium channels; lengthens cardiac potential. Slows cardiac conduction through the AV node. Antisymphathetic action and negative inotropic effects in cardiac nodal tissue. Used for ventricular arrhythmias (ventricular tachycardia/ventricular fibrillation) and some atrial arrhythmias (atrial fibrillation, but takes hours)
<b>ONSET</b>	15 minutes
<b>TIME TO PEAK</b>	1 to 4 hours
<b>DURATION</b>	3 to 6 hours
<b>HALF-LIFE</b>	9-36 hours
<b>METABOLISM</b>	Hepatic

<b>ASPIRIN (ASA)</b>	
<b>CLASS:</b>	Platelet aggregation inhibitor, analgesic, antipyretic and anti-inflammatory
<b>ACTION:</b>	Decreases clotting by inactivating cyclooxygenase, interfering with Thromboxane A2 production within the platelets. Thromboxane A2 also causes arteries to constrict. Reduced morbidity/mortality in adults with C/P from an AMI.
<b>ABSORPTION</b>	Rapid
<b>TIME TO PEAK</b>	1-2 hours
<b>METABOLISM</b>	Hydrolyzed to salicylate (active) in GI mucosa, RBC, synovial fluid and blood. Metabolism of salicylate primarily by the liver.

<b>ATROPINE</b>	
<b>CLASS</b>	Parasympatholytic, anticholinergic
<b>ACTION</b>	Blocks the action of acetylcholine at parasympathetic sites in smooth muscle, secretory glands and the CNS. Results in increased cardiac output and dries secretions.
<b>ONSET</b>	Rapid
<b>HALF-LIFE</b>	2-3 hours
<b>DISTRIBUTION</b>	Widely throughout the body; crosses placenta; trace amounts enter breast milk; crosses blood-brain barrier.
<b>METABOLISM</b>	Hepatic

<b>CALCIUM GLUCONATE 10%</b>	
<b>CLASS</b>	Minerals and electrolytes
<b>ACTION</b>	Calcium protects the myocardium from the deleterious effects of hyperkalemia. It stabilizes the cardiac cell membrane.
<b>ADVERSE REACTION</b>	When given too rapidly can cause hypotension, bradycardia and syncope. If administered IM or extravagates it can cause necrosis/abscess. When given to someone on digoxin this may cause sudden death from ventricular fibrillation.
<b>ADMIN</b>	Slow IV push over 2-3 minutes Incompatible with Sodium Bicarbonate in same IV line.
<b>ONSET</b>	Rapid
<b>DURATION</b>	30 minutes - 2hours
<b>SIDE EFFECTS</b>	Chalky taste, N&V, Dry mouth

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.**DEXAMETHASONE**

<b>CLASS</b>	Adrenocortical steroid
<b>ACTION</b>	Binds to the glucocorticoid receptors inhibiting the release of pro-inflammatory signals through cytokine inhibition, resulting in decreased edema, fibrin deposition, capillary leakage and migration of inflammatory cells.
<b>ONSET</b>	5-15 min(IV); 30 min (PO)60 minutes
<b>DURATION</b>	3 days
<b>HALF-LIFE</b>	4 hours

**DEXTROSE (D50) IN WATER**

<b>CLASS</b>	Carbohydrate
<b>ACTION</b>	Replenishes blood glucose levels, reversing hypoglycemia.
<b>METABOLISM</b>	Metabolized to carbon dioxide and water.

**DIMENHYDRINATE (GRAVOL)**

<b>CLASS</b>	Antiemetic, Antihistamine
<b>ACTION</b>	Competes with histamine for H1-receptor sites on effector cells in the GI tract, blood vessels and respiratory tract; blocks chemoreceptor trigger zone, diminishes vestibular stimulation and depresses function through its central anticholinergic activity.
<b>ONSET</b>	1-5 minutes (IV). 15-30 minutes (oral)
<b>PEAK EFFECTS</b>	1-2 hours
<b>DURATION</b>	3-6 hours

**DIPENHYDRAMINE (BENADRYL)**

<b>CLASS</b>	Antihistamine
<b>ACTION</b>	Competes with histamine and H1-receptor sites on effector cells in the GI tract, blood vessels and respiratory tract; anticholinergic and sedative effects are also seen.
<b>ONSET</b>	1-5 minutes (IV). 1-3 hours (oral)
<b>PEAK EFFECTS</b>	1-2 hours (IV). 2-4 hours (oral)
<b>HALF-LIFE</b>	2-10 hours
<b>DURATION</b>	4-8 hours



DOPAMINE	
<b>CLASS</b>	Sympathomimetic agent
<b>ACTION</b>	Stimulates both adrenergic and dopaminergic receptors, lower doses are mainly dopaminergic stimulating and produce renal and mesenteric vasodilation. Higher doses have both dopaminergic and $\beta$ 1-adrenergic stimulating and produce cardiac stimulation and renal vasodilation. Large doses stimulate $\alpha$ -adrenergic receptors.
<b>ONSET</b>	5 minutes
<b>HALF-LIFE</b>	2 minutes
<b>METABOLISM</b>	Renal, hepatic and plasma (25% gets converted to norepinephrine).

EPINEPHRINE	
<b>CLASS</b>	Sympathomimetic agent
<b>ACTION</b>	Stimulate $\beta$ 1, $\alpha$ 1 and $\beta$ 2-adrenergic receptors resulting in relaxation of smooth muscle of the bronchial tree, cardiac stimulation (increasing myocardial O2 consumption) and dilation of skeletal muscle vasculature. Small doses can cause vasodilation via $\beta$ 2-vascular receptors; large doses may produce constriction of skeletal and vascular smooth muscle.
<b>ONSET</b>	5-10 minutes (bronchodilation).
<b>METABOLISM</b>	Hepatic

FENTANYL	
<b>CLASS</b>	Analgesic, opioid
<b>ACTION</b>	Binds to opioid mu-receptors in the CNS causing inhibition of ascending pain pathways, altering the perception of and response to pain; produces generalized CNS depression, respiratory depression, and can cause apnea. Can cause muscle rigidity if rapid IV injection.
<b>ONSET</b>	IV: almost immediately IN: 5-15 minutes
<b>PEAK EFFECT</b>	IV: 6 minutes IN: 12 minutes
<b>METABOLISM</b>	Hepatic



HYDROCORTISONE	
<b>CLASS</b>	Adrenal glucocorticoid, corticosteroid
<b>ACTION</b>	Short-acting corticosteroid; when used in adrenal crisis or adrenocortical deficiency it replaces/mimics the person's own cortisol which regulates glucose, regulates the immune system, and is released during stressors to help support the cardiovascular system
<b>ONSET</b>	1-2 hours
<b>PEAK EFFECT</b>	1.5 – 2 hours
<b>DURATION</b>	6-12 hours
<b>METABOLISM</b>	Hepatic

HYDROMORPHONE	
<b>CLASS</b>	Opioid analgesic
<b>ACTION</b>	Binds to the mu-opioid receptors in the CNS causing inhibition of the ascending pain pathways, altering the perception of and response to pain. Produces generalized CNS depression
<b>ONSET</b>	5 minutes
<b>DURATION</b>	3-4 hours
<b>HALF-LIFE</b>	2-3 hours

IBUPROFEN	
<b>CLASS</b>	Antipyretic, analgesia and non-steroid anti-inflammatory
<b>ACTION</b>	Its pharmacological effects are believed to be due to inhibition COX-2 which decreases the synthesis of prostaglandins involved in mediating inflammation, pain, fever and swelling. Antipyretic effects may be due to action on the hypothalamus, resulting in an increased peripheral blood flow, vasodilation, and subsequent heat dissipation.
<b>PEAK EFFECT</b>	120 minutes
<b>ONSET</b>	15 minutes
<b>DURATION</b>	4-6 hours
<b>ADVERSE EFFECTS</b>	HTN, MI, GI bleeding, increased the risk of gastric ulcers and damage and renal failure.
<b>METABOLISM</b>	Ibuprofen and its metabolites pass easily across the placenta. More than 90% of an ingested dose is excreted in the urine as metabolites or their conjugates.

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.**KETOROLAC (TORADOL)**

<b>CLASS</b>	Analgesic, antipyretic and non-steroid anti-inflammatory
<b>ACTION</b>	Blocks prostaglandin formation thereby decreasing nociceptor stimulation.
<b>ONSET</b>	10 minutes (IM/IV)
<b>PEAK EFFECT</b>	2-3 hours
<b>DURATION</b>	6-8 hours
<b>METABOLISM</b>	Mostly the hepatic

**LIDOCAINE (XYLOCAINE)**

<b>CLASS</b>	Class 1b antiarrhythmic
<b>ACTION</b>	Suppresses automaticity of conductive tissue by increasing the electrical stimulus threshold of the ventricles, His-Purkinje system and spontaneous depolarization of the ventricles during diastole (by direct action on the tissues). Blocks both the initiation and conduction of nerve impulses by decreasing the neural membranes permeability to Na ions, which results in inhibition of depolarization with resultant blockade of conduction.
<b>ONSET</b>	45-90 seconds
<b>DURATION</b>	10-20 minutes
<b>METABOLISM</b>	90% hepatic

**MIDAZOLAM (VERSED)**

<b>CLASS</b>	Benzodiazepine, CNS depressant, Sedative and Amnesic
<b>ACTION</b>	Binds to stereospecific benzodiazepine receptors on the post-synaptic GABA neuron at several sites within the CNS (including limbic system and reticular formation). Enhancement of the inhibitory effect of GABA on neural excitability results by increased neural membrane permeability to chloride ions. This shift in chloride.
<b>ONSET</b>	45-90 seconds
<b>DURATION</b>	10-20 minutes
<b>METABOLISM</b>	90% hepatic

MORPHINE	
<b>CLASS</b>	Opioid analgesia
<b>ACTION</b>	Binds to opiate receptors in the CNS causing inhibition of ascending pain pathways, altering the perception of and response to pain; produces generalized CNS depression.
<b>ONSET</b>	2-5 minutes (IV)
<b>PEAK EFFECT</b>	20 minutes (IV)
<b>METABOLISM</b>	Hepatic

Airway /  
Breath.Cardiac /  
Circula.

NALOXONE (NARCAN)	
<b>CLASS</b>	Narcotic Antagonist
<b>ACTION</b>	Competitive narcotic antagonist. Displaces and narcotics bound to opiate receptor sites reversing their effects.
<b>ONSET</b>	2-5 minutes (IM). 8-18 minutes (IN). 2 minutes (IV)
<b>HALF-LIFE</b>	3-4 hours (neonates). 0.5-1.5 hours (adults)
<b>DURATION</b>	30-120 minutes
<b>DISTRIBUTION</b>	Crosses placenta
<b>METABOLISM</b>	Hepatic

LOC

Pain/  
Sed./  
Nausea

Proced.

NITROGLYCERIN	
<b>CLASS</b>	Coronary vasodilator, smooth muscle relaxant and anti-anginal
<b>ACTION</b>	Vasodilation of peripheral veins and arteries with more prominent effects on the veins. Reduces myocardial oxygen demand by decreasing preload; may modestly reduce afterload; dilates coronary arteries and improves collateral flow to ischemic tissues. In smooth muscle, nitric oxide activates smooth muscle relaxation.
<b>ONSET</b>	1-3 minutes (SL). 15-30 minutes (topical). 30 minutes (transdermal)
<b>HALF-LIFE</b>	1-4 minutes
<b>DURATION</b>	25 minutes (SL), 7 hours (topical), 10-12 hours (transdermal)
<b>METABOLISM</b>	Extensive first-pass effect; hepatic, RBC and vascular walls

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.**ONDANSETRON**

<b>CLASS</b>	5-HT3 antagonist
<b>ACTION</b>	Selective 5-HT3 receptor antagonist. Mechanism of action through blocking the action of 5-HT3 selectively peripherally and through the vagus nerve, a natural substance that may cause nausea and vomiting. Centrally the chemoreceptor trigger zone is effected.
<b>ONSET</b>	20-30 min
<b>HALF-LIFE</b>	3-6 hrs (PO); 5-8 HRS (IV, IM)
<b>DURATION</b>	4-8 hrs (PO); 5-8 hrs (IV, IM)

**OXYTOCIN**

<b>CLASS</b>	Hormone
<b>ACTION</b>	Promotes uterine contractions by increasing intracellular calcium levels. Greatest effect during labor at term due to increased oxytocin receptor concentrations in uterine myometrial tissue
<b>ONSET</b>	3-5 min
<b>HALF-LIFE</b>	2-3 hrs
<b>DURATION</b>	1-6 min

**SALBUTAMOL (VENTOLIN)**

<b>CLASS</b>	Sympathomimetic, $\beta_2$ agonist
<b>ACTION</b>	Relaxes bronchial smooth muscle by action on $\beta_2$ -receptors with little effect on heart rate
<b>ONSET</b>	10 minutes (Neb/Inhalation)
<b>HALF-LIFE</b>	3-8 hours (inhaled)
<b>DURATION</b>	3-4 hours (Neb/Inhalation)
<b>METABOLISM</b>	Hepatic to an inactive sulfate

XYLOMETAZOLINE (OTRIVIN)	
<b>CLASS</b>	Sympathomimetic Adrenergic Alpha-agonist, decongestant
<b>ACTION</b>	When sprayed into the nares, causes vasoconstriction of the nasal mucosa, resulting in a decrease in blood flow in the nasal passages, decreased nasal congestion, and may help stop epistaxis.
<b>ONSET</b>	5-10 minutes

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

Destinat.  
Guide.

Intentionally Left Blank





## Contact

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



## Physician On-Scene Reference

### For the Paramedic:

If a paramedic encounters a physician on-scene that would like to assist or direct care, the paramedic will follow the Ontario BLS-PCS for any BLS care and the Medical Directives in this document for any ALS care. Inform the physician that paramedics are not able to accept delegation for controlled medical acts from any physician other than an affiliated Base Hospital Physician. The paramedic may show the following information to the on-scene physician to assist in next steps and provide further information.

### To the On-Scene Physician:

Thank you for your assistance.

The paramedics would usually take responsibility for the patient(s) upon their arrival. If, as a physician, you wish to assist with the emergency after the providers have arrived you have three options:

1. Offer your assistance or suggestions that follow the Ontario Basic Life Support Patient Care Standards and/or the Paramedic Medical Directives. If your instructions are not in accordance with these documents, the paramedics cannot follow this direction but can contact the Regional Base Hospital Physician for direction.
2. Take complete responsibility for patient - in which case you will need to accompany the patient to hospital. The paramedics will assist you, but cannot perform skills that do not follow their directives. You may be asked to show identification that you are a physician licensed to practice medicine in Ontario.
3. Request to speak with the Regional Base Hospital Physician (via patch) to offer advice and consult on the best management of the patient(s).

OMC Physician Email Addresses and CPSO Numbers (April 30, 2025)

	Physician	RBH	CPSO #
1	Michael Austin	RPPEO	93761
2	Kim Barker	CPER	72299
3	Dan Beamish	RPPEO	107026
4	Melissa Bouwsema	RPPEO	115439
5	Renee Bradley	RPPEO	112019
6	Nick Costain	RPPEO	97795
7	Richard Dionne	RPPEO	75776
8	Anthony Dixon	CPER	080763
9	Chris Evans	RPPEO	86337
10	Natasha Ewert	HSN	103531
11	Mark Froats	RPPEO	90680
12	Justin Godbout	RPPEO	108994
13	Jonathan Gravel	RPPEO	112054
14	Amy McCulloch	CPER	95286
15	Paul Miller	CPER	76090
16	Simeon Mitchell	RPPEO	101098
17	Jennifer Moore	HSN	98157
18	Sara-Pier Piscopo	RPPEO	112007
19	Jason Prpic	HSN	73974
20	Rupinder Sahsi	CPER	76338
21	Catherine Sellens	CPER	67935
22	Paula Sneath	CPER	115152
23	Ayesha Zia	RPPEO	100347
24	Connor Inglis	RPPEO	118659
25	Louis Gascon-Tetreault (PTM Fellow until July 1)	RPPEO	155640
26	Abdulrahman Ahmed Alkadhail (PTM Fellow until July 1)	RPPEO	147484
27	Margaret Vincent (PTM Fellow until December 31)	RPPEO	120824
28	Ian Buchanan	CPER	95889

## Contact Information

430 McNeilly Road, Unit 201  
Stoney Creek, Ontario L8E 5E3  
Telephone Number: 905-521-2100 x71223  
Fax Number: 905-643-1104

Name:	Position:	EXT:	Mobile:	Email Address:
<b>Tim Dodd</b>	Regional Program Manager/ Director		905-515-4818	tdodd@cper.ca
<b>Dr. Paul Miller</b>	Regional Medical Director			millerpa@hhsc.ca
<b>Dr. Rupinder Sahsi</b>	Assistant Medical Director			rupinder@sahsi.net
<b>Dr. Ian Buchanan</b>	Assistant Medical Director			ian.buchanan@medportal.ca
<b>Dr. Gina Agarwal</b>	Senior Medical Advisor			agarg@mcmaster.ca
<b>Colette Easton</b>	Administration Assistant (To the Directors)	71226		ceaston@cper.ca
<b>Audrey Collie</b>	Administration Assistant (To the Programs)	71229		acollie@cper.ca
<b>Jackie Swing</b>	Administration Assistant	71223		jswing@cper.ca
<b>Angela Burgess</b>	Lead Quality Specialist		289-286-0975	aburgess@cper.ca
<b>Kailash Selvaratinam</b>	Quality Specialist		905-870-4457	kselvar@cper.ca
<b>Carrie Schneider</b>	Quality Specialist		519-503-6632	cschneider@cper.ca
<b>Kathy Winter</b>	Quality Specialist		416-436-5428	winterkat@hhsc.ca
<b>Stephanie Coletta</b>	Lead Paramedic Educator		905-515-0659	scoletta@cper.ca
<b>David Plyley</b>	Paramedic Educator		289-219-1952	dplyley@cper.ca
<b>Bhaven Kapadia</b>	Paramedic Educator			bkapadia@cper.ca
<b>Cyprian Czechowski</b>	Paramedic Educator		647-992-1695	cczechowski@cper.ca
<b>Katie Turcotte</b>	Data Analyst		905-977-1739	kturcotte@cper.ca
<b>Peggy D'Eath</b>	Outreach Specialist		365-324-8389	pdeath@cper.ca

## HHS Centre for Paramedic Education and Research Additional Contact Information Reference

### Central Ambulance Communication Centres (CACC):

CACC – Cambridge	800-265-2215
CACC – Hamilton	905-574-1414
CACC – Hamilton (Alternate)	800-263-5767
CACC – Niagara Ambulance Communication Centre	905-704-4005 866-895-6227

### Emergency Medical Services:

Brant / Brantford Paramedic Service	519-756-4570
Dufferin County Paramedic Service	519-941-9608
Guelph-Wellington Paramedic Service	519-824-1677
Haldimand County Paramedic Services	905-318-5932
Hamilton Paramedic Service	905-546-2424
Niagara EMS	905-641-0827
Norfolk County Paramedic Services	519-426-4115
Region of Waterloo Paramedic Service	519-650-8295
Six Nations Paramedic Services	519-445-4000

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Destinat.  
Guide.

## Community Support Referral Contact Information

The following contact information is provided for cases where:

- ▶ Patients are **refusing** transport to the hospital, and
- ▶ An assessment shows that the patient has the **capacity to refuse**, and
- ▶ The patient does not appear to be of **immediate danger to themselves or others**, and
- ▶ Paramedics have **ongoing concerns** regarding the living conditions in their home (CCAC), their need for victim's support services (victim's services) or the patient's mental health (COAST, Hamilton only)
- ▶ OR the family of a patient needs support services (Victims Services).

These community service organizations are available to assist people with these concerns. Paramedics can give the information directly to the patient or assist them by making the referral on their behalf. Please note that if the Paramedic assists the patient by calling the organization he/she must get the patient's consent to do so. If the Paramedic contacts the organization directly, the agency will require the patient's name, address, phone number and nature of the concern. The Paramedic must then leave the information about the organization called with the patient.



Connecting you with care  
Votre lien aux soins

**CCAC CASC**  
Community  
Care Access  
Centre Centre d'accès  
aux soins  
communautaires

**CCAC (Community Care Access Centre):** provides services for persons with living condition concerns (message can be left).

Brantford CCAC:	800-810-0000
Dufferin County CCAC:	519-925-5452
Guelph-Wellington CCAC:	519-823-2550
Haldimand / Hamilton CCAC:	800-810-0000
Niagara Region CCAC:	800-810-0000
Norfolk / Simcoe CCAC:	800-810-0000
Six Nations (Ohsweken)	519-445-2418
Waterloo - Kitchener CCAC:	519-748-2222

**Victims Services:** provides short-term emotional support and community referral and assistance to victims of crime, tragic circumstance or disaster (24/7).

Brantford	519-752-3140
Cambridge	519-585-2369 / 519-570-5143
Dufferin County	519-942-1452
Guelph-Wellington	519-824-1212 ext. 7304
Haldimand County	800-264-6671
Hamilton Victim Services	905-546-4904
Kitchener	519-585-2369 / 519-570-5143
Niagara Region	905-682-2626
Norfolk County	800-264-6671
Six Nations (Ohsweken)	519-752-3140
Waterloo Region	519-585-2369 / 519-570-5143



**COAST (Crisis Outreach And Support Team):** provides services for persons with mental health concerns in the Hamilton area only (24/7).

Hamilton – Only (24/7) 905 972-8338

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

## Child in Need of Protection

Paramedics have a duty to report under the Child and Family Services Act (CFSA) and this extends to any child they encounter in their professional duties and is not limited to the person (s) requesting 9-1-1 services<sup>1</sup>. This duty overrides any other provincial statute, including any provisions that would otherwise prohibit someone from making a disclosure (i.e. PHIPA). This failure to report a suspicion in the circumstances set out in the CFSA is an offence under the Act.<sup>2</sup>

### Children's Aid Societies in Ontario

#### Dufferin Child and Family Protection Services

Bus: (519) 941-1530

#### Family & Children's Services of Guelph and Wellington County

Bus: (519) 824-2410

#### Children's Aid Society of Hamilton

Bus: (905) 522-1121

#### Catholic Children's Aid Society of Hamilton

Bus: (905) 525-2012

#### Family & Children's Services Niagara

Bus: (888) 937-7731

#### Children's Aid Society of Haldimand and Norfolk

Bus: (519) 587-5437  
Toll Free: (888) 227-5437

#### Brant Family and Children's Services

Bus: (519) 753-8681  
Toll Free: (888) 753-8681

#### Family & Children's Services of the Waterloo Region

Bus: (519) 576-0540

<sup>1</sup> Training Bulletin 116 -Child in Need of Protection Standard March 2015 Version 1.0

<sup>2</sup> Basic Life Support Patient Care Standards –Version 2.2





# Destination Guidelines

ADVANCED CARE PARAMEDIC MEDICAL DIRECTIVES



# Field Trauma Triage Standards

## Definitions

For the purposes of the *Field Trauma Triage Standard*:

### Regionally Designated Equivalent Hospital

means an appropriately resourced hospital facility as defined by the Regional Trauma Network of Critical Care Services Ontario and included in a local PPS.

### Transport Time

means the time from scene departure to time of arrival at destination.

## General Directive

The paramedic shall follow the procedure below when conducting field triage of patients injured by a traumatic mechanism or who show evidence of trauma.

The paramedic shall also use this standard to assess the clinical criteria (*i.e.* to determine if the patient meets the clinical criteria) as required by the *Air Ambulance Utilization Standard*.

The paramedic shall consider using the Trauma Termination of Resuscitation (TOR) contained in the *Trauma Cardiac Arrest Medical Directive* as per the ALS PCS.

CACC/ACS may authorize the transport once notified of the patient's need for re-direct or transport under the *Field Trauma Triage Standard*.

## Procedure

The paramedic shall:


1. assess the patient to determine if he/she has one or more of the following **physiological criteria** (Step 1):
  - a. Patient does not follow commands,
  - b. Systolic blood pressure <90mmHg, or
  - c. Respiratory rate <10 or ≥30 breaths per minute or need for ventilatory support (<20 in infant aged <1 year);
2. if the patient meets the physiological criteria listed in paragraph 1 above, **AND** the land transport time is estimated to be <30 minutes\* to a Lead Trauma Hospital (LTH) or regionally designated equivalent hospital, transport the patient directly to the LTH or regionally designated equivalent hospital;
3. if the patient does not meet the criteria listed in paragraphs 1 and 2, assess the patient to determine if he/she has one or more of the following **anatomical criteria** (Step 2):

	Intro
	Airway / Breath.
<ul style="list-style-type: none"> <li>a. Any penetrating injuries to head, neck, torso and extremities proximal to elbow or knee,</li> <li>b. Chest wall instability or deformity (e.g. flail chest),</li> <li>c. Two or more proximal long-bone fractures,</li> <li>d. Crushed, de-gloved, mangled or pulseless extremity,</li> <li>e. Amputation proximal to wrist or ankle,</li> <li>f. Pelvic fractures,</li> <li>g. Open or depressed skull fracture, or</li> <li>h. Paralysis;</li> </ul>	Cardiac / Circula.
4. if the patient meets the anatomical criteria listed in paragraph 3 above and the land transport time is estimated to be <30 minutes* to the LTH or regionally designated equivalent hospital, transport the patient directly to the LTH or regionally designated equivalent hospital;	LOC
5. if unable to secure the patient's airway or survival to the LTH or regionally designated equivalent hospital is unlikely, transport the patient to the closest emergency department despite paragraphs 2 and 4 above;	Pain/ Sed./ Nausea
6. despite paragraph 5 above, transport the patient directly to an LTH or regionally designated equivalent hospital if the patient has a penetrating trauma to the torso or head/neck, and meets <b>ALL</b> of the following: <ul style="list-style-type: none"> <li>a. Vital signs absent yet not subject to TOR described in the <i>General Directive</i> above, and</li> <li>b. Land transport to the LTH or regionally designated equivalent hospital is estimated to be &lt;30 minutes*;</li> </ul>	Proced.
7. if the patient does not meet the physiological or anatomical criteria listed above, use the following <b>criteria</b> to determine if the patient may require other support services at the LTH or regionally designated equivalent hospital as a result of his/her traumatic <b>mechanism of injury</b> (Step 3): <ul style="list-style-type: none"> <li>a. Falls <ul style="list-style-type: none"> <li>i. Adults: falls <math>\geq 6</math> metres (one story is equal to 3 metres)</li> <li>ii. Children (age &lt;15): falls <math>\geq 3</math> metres or two to three times the height of the child</li> </ul> </li> <li>b. High Risk Auto Crash <ul style="list-style-type: none"> <li>i. Intrusion <math>\geq 0.3</math> metres occupant site; <math>\geq 0.5</math> metres any site, including the roof</li> <li>ii. Ejection (partial or complete) from automobile</li> <li>iii. Death in the same passenger compartment</li> <li>iv. Vehicle telemetry data consistent with high risk injury (if available)</li> </ul> </li> <li>c. Pedestrian or bicyclist thrown, run over or struck with significant impact (<math>\geq 30</math> km/hr) by an automobile</li> <li>d. Motorcycle crash <math>\geq 30</math> km/hr;</li> </ul>	Pall Care / Research
	Chemical Exposure
	Medical Refer.
8. if the patient meets the mechanism of injury criteria listed in paragraph 7 above, <b>AND</b> the land transport time is estimated to be <30 minutes* to an LTH or regionally designated equivalent hospital, determine the need for patient transport to the LTH or regionally designated equivalent hospital;	Medic. Info.
	Contact
Destination Guidelines    Field Trauma Triage Standards BLS PCS 3.4	
	238
	Destinat. Guide.

Intro	
Airway / Breath.	
Cardiac / Circula.	
LOC	<p>9. in conjunction with the physiological, anatomical, and mechanism of injury criteria listed above, consider the following <b>special criteria</b> (Step 4):</p> <ol style="list-style-type: none"> <li>Age <ol style="list-style-type: none"> <li>Risk of injury/death increases after age 55</li> <li>SBP &lt;110 may represent shock after age 65</li> </ol> </li> <li>Anticoagulation and bleeding disorders</li> <li>Burns <ol style="list-style-type: none"> <li>With trauma mechanism: triage to LTH</li> </ol> </li> <li>Pregnancy ≥20 weeks; and</li> </ol> <p>10. if the patient meets any of the special criteria listed above, <b>AND</b> the land transport time is estimated to be &lt;30 minutes* to an LTH or regionally designated equivalent hospital, determine the need for patient transport to the LTH or regionally designated equivalent hospital.</p> <p><b>*Note: The 30 minute transport time may be amended to up to 60 minutes as per an ambulance service PPS, but may not exceed 60 minutes.</b></p>
Pain/ Sed./ Nausea	
Proced.	
Pall Care / Research	
Chemical Exposure	
Medical Refer.	
Medic. Info.	
Contact	
Destinat. Guide.	<p>Destination Guidelines</p> <p>239</p>

# Paramedic Prompt Card for Field Trauma Triage Standard

This prompt card provides a quick reference of the *Field Trauma Triage Standard* contained in the *Basic Life Support Patient Care Standards* (BLS PCS). Please refer to the BLS PCS for the full standard.



\*The 30 minute transport time may be amended to up to 60 minutes as per an ambulance service PPS, but may not exceed 60 minutes.

\*\*If unable to secure the patient's airway or survival to the LTH or regionally designated equivalent hospital is unlikely, transport the patient to the closest ED (unless patient has penetrating trauma to the torso or head/neck). Consider the Trauma TOR as per the ALS PCS.

# Air Ambulance Utilization Standard

## General Directive

Requests for an on-scene air ambulance response should meet at least one of the bulleted operational criteria **PLUS** one of the clinical criteria (*e.g.* known clinical criteria as listed in the *Field Trauma Triage Standard* or from the bulleted list of medical or obstetrical criteria listed below).

## Procedure

### The paramedic shall:

1. assess the scene response to meet one or more of the following **operational criteria**:
  - a. The land ambulance is estimated to require more than 30 minutes to reach the scene and the air ambulance can reach the scene quicker.
  - b. The land ambulance is estimated to require more than 30 minutes to travel from the scene to the closest appropriate hospital\* and the air ambulance helicopter can reach the scene and transport the patient to the closest appropriate hospital\* quicker than the land ambulance.
  - c. The estimated response for both land and air is estimated to be greater than 30 minutes, but approximately equal, and the patient needs care which cannot be provided by the responding land ambulance.
  - d. There are multiple patients who meet the clinical criteria and the local land ambulance resources are already being fully utilized.
2. if the scene response meets the requirements of paragraph 1 above, assess the patient to determine if he/she meets one or more of the following **clinical criteria**:
  - a. Patients meeting the criteria listed in the *Field Trauma Triage Standard*.
  - b. Patients meeting one or more of the following:
    - i. **Medical**:
      1. Shock, especially hypotension with altered mentation (*e.g.* suspected aortic aneurysm rupture, massive gastrointestinal bleed, severe sepsis, anaphylaxis, cardiogenic shock, *etc.*)
      2. Acute stroke with a clearly determined time of onset or last known to be normal <6.0 hours
      3. Altered level of consciousness (GCS <10)
      4. Acute respiratory failure or distress
      5. Suspected STEMI or potentially lethal dysrhythmia
      6. Resuscitation from respiratory or cardiac arrest
      7. Status epilepticus
      8. Unstable airway or partial airway obstruction

	Intro
ii. <b>Obstetrical:</b>	
1. Active labour with abnormal presentation ( <i>i.e.</i> shoulder, breech or limb)	Airway / Breath.
2. Multiple gestation and active labour	
3. Umbilical cord prolapse	Cardiac / Circula.
4. Significant vaginal bleeding (suspected placental abruption or placenta previa or ectopic pregnancy);	
3. in conjunction with the ACO, assess if an on-scene air ambulance helicopter is appropriate, based on:	
a. the perceived severity of the reported injuries and without confirmation that the clinical criteria have been met, or	
b. the patient cannot reasonably be reached by land ambulance ( <i>e.g.</i> sites without road access such as islands; geographically isolated places, <i>etc.</i> );	LOC
4. if the requirements listed in paragraph 2 or 3 above are met, request an on-scene air ambulance helicopter response:	
a. Provide the ACO with the information set out in operational and clinical criteria above. In order for the ACO to determine if an air ambulance response and transport will be quicker than land ambulance, the paramedic will provide the ACO with the estimated time to prepare the patient for transport, identify separately any time required for patient extrication, provide the estimated land ambulance driving time to the closest appropriate hospital and any additional information as required.	Pain/ Sed./ Nausea
b. The paramedics shall not delay patient transport by waiting for the air ambulance helicopter, unless the air ambulance helicopter can be seen on its final approach to the scene. If the air ambulance helicopter is en route but not on final approach to the scene, and the land paramedics have the patient in his/her ambulance, then the land ambulance will proceed to the closest local hospital with an emergency department. The air ambulance helicopter will proceed to that local hospital and, if appropriate, assist hospital personnel prepare the patient for rapid evacuation.	Proced.
c. While en route to the local hospital, paramedics may rendezvous with the air ambulance helicopter if:	Pall Care / Research
i. the air ambulance helicopter is able to land along the direct route of the land ambulance; and	
ii. it would result in a significant reduction in transport time to the most appropriate hospital.	Chemical Exposure
5. if the call's circumstances and patient(s) fail to meet the criteria set out in this standard and an air ambulance helicopter is known to be responding based on the merits of the initial request for ambulance service, contact the CACC/ACS and advise that an on-scene air ambulance helicopter response is not required and why it is not required.	Medical Refer.
	Medic. Info.
	Contact

---

**Guideline****Air Ambulance Helicopter Landing Site Safety and Coordination**


Upon confirmation that the air ambulance helicopter is responding, the paramedic shall follow the guidelines set out by the Ornge Aviation Safety Department, which can be found on Ornge's "Aircraft Safety" website at: <https://www.ornge.ca/aircraft-safety>.

**Other Use of Air Ambulance Helicopter**

- Air ambulance helicopters are not permitted to respond to night calls which require a landing at a site other than night licensed airports, helipads or night approved remote landing sites.
  - Air ambulance helicopters are not permitted to conduct search and rescue calls.
  - In cases where a land ambulance can reach the patient(s) and an on-scene response by air ambulance helicopter is appropriate, the ACO will assign a land ambulance and continue the land response until the flight crew requests that the land ambulance be cancelled.
  - In cases where a land ambulance arrives on-scene prior to the air ambulance helicopter, paramedics shall inform the CACC/ACS as clinical events occur.
-



# Deceased Patient Standards



## Deceased patient means a patient who is:

- Obviously dead – code 5
- Subject to a MCOB presented to the paramedic
- VSA and subject to a valid DNR
- VSA and is subject to a Termination of Resuscitation Order
- VSA and is subject to a Withdrawal Resuscitation Order

**Note:** When a Termination of resuscitation Order is received, and the deceased person has not been removed from the place of death, paramedics should not remove the body, but rather they should follow the appropriate procedure as outlined.

## Paramedic Prompt Card for Acute Stroke Bypass Protocol

This prompt card provides a quick reference of the *Acute Stroke Protocol* contained in the *Basic Life Support Patient Care Standards (BLS PCS)*.

### Indications under the Acute Stroke Protocol

Redirect or transport to the closest or most appropriate Designated Stroke Centre (DSC)\* will be considered for patients who meet **BOTH** of the following:

1. Present with a new onset of at least one of the following symptoms suggestive of the onset of an acute stroke:
  - a. Unilateral arm/leg weakness or drift.
  - b. Slurred speech or inappropriate words or mute.
  - c. Unilateral facial droop.
2. Can be transported to arrive at a Designated Stroke Centre within 6 hours of a clearly determined time of symptom onset or the time the patient was last seen in a usual state of health.

**Inform the CACC/ACS to aid in the determination of the most appropriate destination.**

\*A Regional Stroke Centre, District Stroke Centre or Telestroke Centre regardless of EVT capability.

### Large Vessel Occlusion (LVO) Assessment

Perform a secondary screen for LVO stroke using the Los Angeles Motor Scale (LAMS) for all probable stroke patients presenting within 24 hours of stroke symptom onset.

- a. if LAMS is greater than or equal to 4 ( $\geq 4$ ), classify the patient as CTAS 2
- b. inform the receiving hospital whether "LVO Clinical Screen is positive or negative" \*\*

\*\* In select regions, LVO Clinical Screen + patients, presenting within 6 hours of stroke symptom onset, may be redirected to the closest EVT centre.

### Contraindications under the Acute Stroke Protocol

**ANY** of the following exclude a patient from being transported under the Acute Stroke Protocol:


1. CTAS Level 1 and/or uncorrected airway, breathing or circulatory problem.
2. Symptoms of the stroke resolved prior to paramedic arrival or assessment\*\*\*.
3. Blood sugar  $< 3$  mmol/L\*\*\*.
4. Seizure at onset of symptoms or observed by paramedics.
5. Glasgow Coma Scale  $< 10$ .
6. Terminally ill or palliative care patient.
7. Duration of out of hospital transport will exceed two hours.

**\*\*\*Patients whose symptoms improve significantly or resolve during transport will continue to be transported to a Designated Stroke Centre.**

\*\*\*If symptoms persist after correction of blood glucose level, the patient is not contraindicated.

**CACC/ACS will authorize the transport once notified of the patient's need for redirect or transport under the Acute Stroke Protocol.**

## Reporting to FACS Niagara



Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

## Paramedic Prompt Card for Sepsis



### Paramedic Prompt Card for Sepsis Reference

#### Suspected or Confirmed Signs and Symptoms of Infection?

**Skin:** Cellulitis, Wound, Burns

**Immunocompromised Neuro:** LOC changes, Weakness,  
Indwelling Medical Device

**Chest:** Cough, SOB, Recent Surgery/Invasive Procedure

**Abdomen:** Pain, Vomiting, Diarrhea, History of Fever or Rigors  
(shakes)

**Urine:** Dysuria, Frequency, Odour

**Age :**  $\geq 18$

#### At Least 2 OR MORE:

**Temperature:**  $< 36^{\circ}\text{C}$  OR  $\geq 38^{\circ}\text{C}$

**Pulse:**  $\geq 90$  bpm

**Respiratory Rate:**  $\geq 20$  bpm

#### And at least ONE of the following

Signs of Hypoperfusion ( $\text{O}_2$  Sat  $< 92\%$ )

Systolic BP  $< 90$  mmHg

New Altered mental status

#### Suggested Treatment

IV access obtained

Intravenous & Fluid Therapy Directive (bolus)

Notify ED of **\*Sepsis Alert\***

YES

NO

# Paramedic Prompt Card for Sepsis (NEMS)



## Paramedic Prompt Card for Sepsis Reference

	YES	NO
<b><u>Suspected or Confirmed Signs and Symptoms of Infection?</u></b> <ul style="list-style-type: none"> <li>▶ <b>Skin:</b> Cellulitis, Wound, Burns</li> <li>▶ <b>Immunocompromised /Neuro:</b> LOA changes, Weakness, Indwelling Medical Device , Chemotherapy</li> <li>▶ <b>Chest:</b> Cough, SOB, Recent Surgery/Invasive Procedure</li> <li>▶ <b>Abdomen:</b> Pain, Vomiting, Diarrhea with a history of fever or rigors</li> <li>▶ <b>Urine:</b> Dysuria, Frequency (increased or decreased), Odour</li> </ul>		
<b>Age :</b> ≥ 18		
<b><u>At Least 2 OR MORE of the following:</u></b> <ul style="list-style-type: none"> <li>▶ <b>Temperature:</b> &lt; 36° C OR ≥ 38° C</li> <li>▶ <b>Pulse:</b> ≥ 90 bpm</li> <li>▶ <b>Respiratory Rate:</b> ≥ 20bpm</li> </ul>		
<b><u>And at least ONE of the following</u></b> <ul style="list-style-type: none"> <li>▶ Signs of Hypoperfusion (mottled extremities, poor cap refill, etc)</li> <li>▶ Systolic BP &lt;90mmHg</li> <li>▶ New altered LOA</li> </ul>		
If you answer yes to all of the above then Notify ED of <b>*Sepsis Alert*</b>		
<b><u>Suggested Treatment</u></b> <ul style="list-style-type: none"> <li>▶ IV access</li> <li>▶ Intravenous &amp; Fluid Therapy Directive</li> <li>▶ If the patient clearly meets the Sepsis Alert <b>AND</b> they do not meet the Medical Directive for fluid therapy, consider contacting the BHP for IV fluid orders.</li> </ul>		

Intro

Airway /  
Breath.Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

## Niagara EMS Hospital Destination Policy



### Policy # IV 3.12a Hospital Destination Policy

July 5, 2023

#### HOSPITAL DESTINATION POLICY - Niagara Region

##### The Paramedic will:

Make a decision regarding receiving facility and transport the patient to that facility or an alternate facility as confirmed or directed by:

- an ambulance dispatcher, or
- an attending physician, with dispatch confirmation, or
- a base hospital physician, with dispatch confirmation, or
- approved local transfer guidelines, or
- the patient, with dispatch approval.

**In the absence of direction, transport to the closest or most appropriate hospital emergency department capable of providing the medical care apparently required by the patient. The goal is to expedite time to definitive care. When there are two or more hospitals equal in time from the level 1 or 2 patient, the Paramedic may choose among available sites in consultation with NEMS Communications.**

If in the paramedic's judgment, the patient can be managed en route the patient will be transported to the most appropriate hospital (as indicated below).

If the patient deteriorates during transport, and survival to the directed receiving facility is questionable, **the paramedic will** transport the patient to the **closest** or most appropriate hospital emergency department capable of providing the medical care immediately required by the patient. **The paramedic will** immediately notify dispatch of any destination change, and notify or ask dispatch to **notify** the **initial** and **receiving** facility.

Patient preference for a specific hospital, other than the closest, will be considered where resources permit based on clinical factors or continuity of care.

CONDITION	DESCRIPTION	DESTINATION
<b>TRAUMA</b>	<p><u>Paramedics/ Dispatchers will consider the Air Ambulance Utilization Standard for FTT</u></p> <hr/> <p>All trauma patients meeting Field Trauma Triage (FTT) Standard Criteria where the incident location is within 60 minutes transport time to a Lead Trauma Centre will be transported to the Lead Trauma Centre in accordance with the guidelines (Policy IV-3.12h).</p> <p>*If transport time to Lead Trauma Centre will exceed 60 minutes, or survival to Lead Trauma Hospital is unlikely (see exception in Policy IV-3.12h), patients meeting FTT criteria will be transported to the closest Emergency Department.</p>	<b>Trauma Center/ Closest Emergency Department *</b>

**Policy # IV 3.12a Hospital Destination Policy**  
July 5, 2023

<b>HEAD TRAUMA</b>  <i>Hospitals with CT: GNG, SCS, WH Sites and WLMH in Niagara HGH Site in Hamilton</i>	<p>All patients with head trauma &amp; <b>an altered LOC not meeting FTT Standard</b> will be taken to the closest hospital with a <b>functioning CT</b>.</p> <p>If they are in active resuscitation then the patient is to be transported to the closest ED.</p>	<b>Closest Emergency Department with a functioning CT (GNG, SCS, WH, WLMH and HGH)</b>
<b>STROKE EMERGENCIES</b>  <i>Stroke Centers: GNG Site and Hamilton General Hospital</i>  <i>Hospitals with CT: GNG, SCG, WH Sites and WLMH in Niagara HGH in Hamilton</i>	<p><b>Patients meeting the criteria of the Paramedic Prompt Card will be taken to the closest Stroke Centre for evaluation (attached)</b></p> <p>Those stroke patients who do <b>not</b> meet the Paramedic Prompt Card criteria will be taken to the closest hospital with a <b>functioning CT</b>.</p> <p>If CT is down at the GNG Site, patients who meet the Provincial Paramedic Prompt Card criteria will be taken to the closest site with a functioning CT with "next on table" priority.</p> <p>They will then be transported to the GNG Site for assessment by the Stroke Team (see attached Appendix A<sub>2</sub> - CT Downtime Contingency Plan for Stroke Thrombolysis (tPA).</p>	<b>Closest Stroke Center</b>
<b>SEXUAL ASSAULT</b>	<p>All victims of sexual assaults will go to the <b>closest</b> hospital for medical clearance.</p> <p>Following patient triage, registration, and physician assessment appropriate transfer arrangements to SCS/HGH will be made by the receiving site if the patient requires sexual assault services.</p>	<b>Closest hospital for medical clearance – then may require transfer to SCS or HGH as appropriate</b>
<b>DIALYSIS EMERGENCIES</b>	<p>All hemo/ peritoneal dialysis with <b>related complaints</b> will be transported to SCS unless the patient is actively being resuscitated, patients will be transported to the closest hospital.</p> <p><b>Consideration will be given to St. Joseph's Health Care Hamilton for patients picked up West of RR24</b></p>	<b>St. Catharines Site or St. Joseph's Health Care</b>
<b>OBSTETRICAL &amp; GYNECOLOGICAL EMERGENCIES</b>	<p>Patients whose chief complaint is Obstetrical in nature will be taken to the SCS (<b>or WLMH if closer</b>) unless active resuscitation is in progress or</p>	<b>St. Catharines Site or WLMH, whichever is</b>

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

**Policy # IV 3.12a Hospital Destination Policy**  
July 5, 2023

	<p>in the case of a laboring patient a presenting fetal part is visible (e.g. crowning). These patients will be taken to the closest Emergency Department.</p> <p>If childbirth has occurred, and no active resuscitation is required, infant and mother should be transported to SCS or <b>WLMH, whichever is closest.</b></p> <p><b>Note: WLMH should typically only be considered for patients greater than 36 weeks gestation.</b></p> <p>Patients whose presentation is highly suggestive of an ectopic pregnancy, for eg. sudden onset severe abdominal pain in a female of child bearing age, should also be considered for transport to SCS or <b>WLMH if closer.</b></p> <p><i>Pregnant patients whose chief complaint is clearly <b>NOT</b> OB/GYN in nature will be transported under the appropriate destination for that complaint as outlined within this policy.</i></p>	<p><b>closest, unless active resuscitation in progress <u>OR</u> presenting fetal part is visible.</b></p>
<b>ONCOLOGY and PALLIATIVE EMERGENCIES</b>	<p>Patients will go to the hospital where they have been receiving treatment within Niagara Region if they can be managed en route.</p> <p><b>Niagara's Regional Cancer Program is the SCS. (Consideration will be given to Juravinski in Hamilton for patients picked up West of RR24)</b></p>	<p><b>St. Catharines Site (consideration for Juravinski West of RR24)</b></p>
<b>PEDIATRIC EMERGENCIES (less than 16 yrs. of age)</b>	<p>Pediatric patients triaged as <b>Level 1, or who require active resuscitation</b>, will go to the closest hospital for immediate assessment and stabilization.</p>	<p><b>If active resuscitation go to closest hospital.</b></p>



**Policy # IV 3.12a Hospital Destination Policy**  
July 5, 2023

	<p>Non-complex Paediatric patients will be taken to the <b>closest</b> hospital.</p> <p><b>Complex patients, such as those with indwelling medical devices, with medically complex histories or injuries, or who are currently receiving treatment at St. Catharines Site, should be transported to the closest hospital with a pediatrician available (SCS in Niagara, MUMC in Hamilton) if the patient can be managed during transport.</b></p> <p>All other patients will be transported to the closest appropriate hospital as outlined in this policy (for example, orthopedics or trauma).</p>	<p>Complex patients go to St. Catharines Site or MUMC depending on location</p>
<b>MENTAL HEALTH EMERGENCIES</b>	<p>Patients of all ages where <b>mental illness is the primary problem</b> will be taken to a <b>schedule 1 facility</b>: SCS in Niagara, or St. Joseph's Healthcare in Hamilton if closer. Patients should be taken to the closest of the two sites.</p> <p>Consideration for previous treatment history with a facility may be considered in choosing an appropriate destination.</p> <p>Patients with a history of mental illness, but in whom the <b>primary problem</b> is medical (i.e. overdose etc.) or surgical emergency will go to the <b>closest</b> appropriate hospital as outlined elsewhere in this policy.</p>	<p><b>If primary problem is medical go to closest hospital.</b></p> <p><b>If Mental Illness is the primary problem then go to St. Catharines Site, or SJHH if closer.</b></p>
<b>ORTHOPEDIC EMERGENCIES</b>	<p>Patients with major orthopedic emergencies (i.e. long bone fracture, spinal or pelvic fracture, open fracture or gross deformity) will be taken to the closest appropriate hospital i.e. where there is an Orthopedic Surgeon on-call if they can be managed en route. <b>This includes HGH to the West.</b> Patients under 16 should be transported to SCS (<b>MUMC if closer</b>)</p> <p><i>Patients with minor orthopedic emergencies (i.e. isolated orthopedic injury, fractured wrist, ankle etc.) will be taken to the closest hospital ED.</i></p>	<p><b>Major: Closest hospital with Ortho (peds to SCS or MUMC)</b></p>

Revised: July 4, 2023

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

**Policy # IV 3.12a Hospital Destination Policy**  
*July 5, 2023*

**PARAMEDIC PROMPT CARD**  
Niagara Regional Acute Stroke Protocol


Refer to **current Paramedic Prompt Card for Acute Stroke Protocol** contained within **the current Basic Life Support Patient Care Standards**.

**The closest Stroke Centre is defined in the CAD.**

**Notify the Receiving Hospital that they will be receiving a "Stroke Alert" patient that meets the Acute Stroke Protocol.**

Transport CTAS Level 2 to the Emergency Department of the closest Stroke Centre.

# Appendix A<sub>2</sub> – CT Downtime Contingency Plan for Stroke Thrombolysis (tPA)



August 9<sup>th</sup>, 2012 - LH

## "The Canadian CSPINE Rule"



The Canadian C-spine Rule is to be utilized only as part of NEMS' "Research Medical Directive to Study the Safety of C-spine Clearance by Paramedics".

### 1. Any ONE High-Risk factor which mandates immobilization?

- Age  $\geq 65$  years
- Dangerous Mechanism\*
- Numbness or tingling in extremities

↓ No

### 2. Any ONE Low-Risk factor which allows safe assessment of range of motion?

- Rear-ended in Simple Rear-end MVC\*\*
- Ambulatory at any time at scene
- No neck pain at scene when asked (answer "yes" if no pain)
- No pain during midline c-spine palpation (answer "yes" if no pain)

↓ Yes

### 3. Patient voluntarily able to Actively Rotate neck 45° left and right when requested, regardless of pain?

↓ Yes

**NO C-SPINE IMMOBILIZATION**

#### Inclusion Criteria

Alert (GCS 15)  
Stable (SBP  $\geq 90$ , RR 10-24)  
Acute Blunt Injury (<48 hrs)

Yes

No

No

**USE C-SPINE  
IMMOBILIZATION**

#### Exclusion Criteria

Boarded & Collared for Other Reasons  
Age < 8  
Penetrating Trauma  
Acute Paralysis  
Known Vertebral Disease  
Referred from another Hospital

#### \*Dangerous Mechanism:

- fall from elevation  $\geq 3$  feet/5 stairs
- axial load to head, e.g. diving
- MVC: rollover, ejection, high speed ( $\geq 100$ km/h)
- motorized recreational vehicles, e.g. ATV, snowmobile
- bicycle collision with object, e.g. post, car

#### \*\*Simple Rearend MVC Excludes:

- pushed into oncoming traffic
- hit by bus/large truck
- rollover
- hit by high speed vehicle ( $\geq 100$ km/h)

# Paramedic Prompt Card for STEMI Hospital Bypass Protocol

This prompt card provides a quick reference of the *STEMI Hospital Bypass Protocol* contained in the *Basic Life Support Patient Care Standards* (BLS PCS). Please refer to the BLS PCS for the full protocol.

## Indications under the STEMI Hospital Bypass Protocol

Transport to a PCI centre will be considered for patients who meet **ALL** of the following:

1.  $\geq 18$  years of age.
2. Chest pain or equivalent consistent with cardiac ischemia/myocardial infarction.
3. Time from onset of current episode of pain  $< 12$  hours.
4. 12-lead ECG indicates an acute AMI/STEMI\*:
  - a. At least 2 mm ST-elevation in leads V1-V3 in at least two contiguous leads; **AND/OR**
  - b. At least 1 mm ST-elevation in at least two other anatomically contiguous leads; **OR**
  - c. 12-lead ECG computer interpretation of STEMI and paramedic agrees.

\*Once activated, continue to follow the STEMI Hospital Bypass Protocol even if the ECG normalizes.

## Contraindications under the STEMI Hospital Bypass Protocol

**ANY** of the following exclude a patient from being transported under the STEMI Hospital Bypass Protocol:

1. CTAS 1 and the paramedic is unable to secure patient's airway or ventilate.
2. 12-lead ECG is consistent with a LBBB, ventricular paced rhythm, or any other STEMI imitator.
3. Transport to a PCI centre  $\geq 60$  minutes from patient contact.\*\*
4. Patient is experiencing a complication requiring PCP diversion:\*\*
  - a. Moderate to severe respiratory distress or use of CPAP.
  - b. Hemodynamic instability or symptomatic SBP  $< 90$  mmHg at any point.
  - c. VSA without ROSC.
5. Patient is experiencing a complication requiring ACP diversion:\*\*
  - a. Ventilation inadequate despite assistance.
  - b. Hemodynamic instability unresponsive/not amenable to ACP treatment/management.
  - c. VSA without ROSC.

\*\*The interventional cardiology program may still permit the transport to the PCI centre.

**CACC/ACS will authorize the transport once notified of the patient's need for bypass under the STEMI Hospital Bypass Protocol.**



# Hamilton Adult Patient Priority System



Airway / Breath.

Cardiac / Circula.

LOC

Pain/ Sed./ Nausea

Proced.

Pall Care / Research

Chemical Exposure

Medical Ref.

Medic. Info.

Contact

Adult patients (≥18 years unless otherwise indicated) will be transported according to applicable MOH Standards. The following clinical presentations will be taken to the facility listed in priority order, as they have the resources to provide the most appropriate medical care. Where applicable, paramedics will utilize "Recognition Tools" to determine if a patient's condition meets a certain criteria.

1

- Pre-arrest
- VSA (except penetrating trauma)
- Unresolved airway compromise

Closest facility\*

2

- Trauma\*\* (≥16 YOA, pregnant patients, VSA with penetrating trauma to head, neck, torso)
- STEMI\*\*
- CVA\*\*
- Major burns >25% BSA or airway problems
- Smoke inhalation injury with altered LOC
- Diving/decompression incidents

HHS - General

3

- Dialysis patient

SJHH

4

- Pregnant patient who are >32 weeks gestation **AND**:
  - In labour **OR**;
  - Expected complications for the mother or fetus\*\*\*\*\*
- Well newborns delivered in the field

1. SJHH - L&D **OR**;  
2. MCH - L&D  
(which ever hospital is the planned site of care)\*\*\*\*\*

5

- Pregnant patient who are 20-31 weeks gestation **AND**:
  - In labour **OR**;
  - Expected complications for the mother or fetus\*\*\*\*\*

MCH - L&amp;D

6

- Pregnant patients who are <20 weeks gestation **AND**:
  - In labour **OR**;
  - Expected complications for the mother or fetus\*\*\*\*\*

1. SJHH - ED planed care-  
**OR**;  
2. Any facility

7

- Suspect or known sexual assault

1. HHS - General **OR**;  
2. HHS - Juravinski

8

- Suspected GI bleed\*\*\*

1. SJHH **OR**;  
2. HHS - Juravinski **OR**;  
3. HHS - General

9

- Suspected hip fracture\*\*\*

1. SJHH **OR**;  
2. HHS - Juravinski **OR**;

10

- Musculoskeletal injury where the paramedic suspects surgery will be required\*\*\*

1. SJHH **OR**;  
2. HHS - General

11

- Psychiatric emergency\*\*\*

SJHH

12


- UCC & SHF patients where arrangements have been made with a receiving ED to continue care

Facility as per direction

13

- Patients that have been an inpatient at a specific hospital within ≤14 days **AND**:
  - Their chief complaint is related to that admission

Facility as per recent history



#### NOTES

\* Patients should be transported to the closest facility unless otherwise directed by provincial guidelines/standards

\*\* As per provincial ALSPCS/BLSPCS Standards, Training Bulletins, or Paramedic Prompt Cards

\*\*\* As per Clinical Recognition Tool

\*\*\*\* When no other PPS criteria are met, Hamilton CACC will equally distribute patients transported to each hospital


\*\*\*\*\* If there has been no prenatal care or the mother is not a patient of MUMC or SIHH, the patient will be taken to either MUMC or SIHH, whichever is closest

\*\*\*\*\* If fetal complications are suspected to be a result of a traumatic or medical emergency involving the mother, paramedics **must transport the mother to an appropriate adult facility first and not MCH**

## Hamilton Pediatric Patient Priority System



Pediatric patients (<18 years unless otherwise indicated) will be transported according to applicable MOH Standards. The following clinical presentations will be taken to the facility listed priority order, as they have the resources to provide the most appropriate medical care. Where applicable, paramedics will utilize "Recognition Tools" to determine if a patient's condition meets a certain criteria.



#### NOTES

\* Patients should be transported to the closest facility unless otherwise directed by provincial guidelines/standards

\*\* As per provincial ALSPCS/BLSPCS Standards, Training Bulletins, or Paramedic Prompt Cards

\*\*\* If fetal complications are suspected to be a result of a traumatic or medical emergency involving the mother, paramedics **must transport the mother to an appropriate facility**

\*\*\*\* As per Clinical Recognition Tool

\*\*\*\*\* If there has been no prenatal care or the mother is not a patient of MUMC or SIHH, the patient will be taken to either MUMC or SIHH, whichever is closest

# GI Bleed Recognition Tool (HPS)



For the purposes of the Patient Priority System:

Patients with possible "GI bleeds" (gastrointestinal bleeding) recognized by the guidelines below should be transported to the appropriate Emergency Department (St. Joseph's Healthcare or HHS Juravinski Site) as directed by CACC.

## INCLUSION

The patient must be;  $\geq 18$  years of age and meet the following:

- 1. Vomiting blood (hematemesis) bright red blood, dark red blood, dark brown/black blood ("coffee grounds") or blood clots.
- 2. Passing red blood rectally (hematochezia) bright red blood, dark red blood or blood clots (with or without stools)
- 3. Passing black stools (melena) sticky, black, "tarry", stools with a typical foul smell – may be mixed with red or maroon blood.

## EXCLUSION

Patients  $< 18$  years should be transported as per the Pediatric Destination Determination Guidelines and not according to this Tool.

Education notes:

Relevant history:

If a patient with a possible "GI bleed" has an extensive history with one site (eg: such as post operative, oncology, dialysis, multiple admissions, or discharged patient), it would be preferable for the patient to be transported to that site (excluding McMaster Children's Hospital or HHS Hamilton General Site).



# Isolated Hip Fracture Recognition Tool (HPS)



For the purposes of the Patient Priority System:

Patients with possible "isolated" hip fracture recognized by the guidelines below should be transported to the Emergency Department as directed by CACC (St. Joseph's Healthcare or HHS Juravinski Site).

## INCLUSION

Mechanism: Fall from sitting (chair), bed, or standing (not height or MVC); may have other minor injuries (i.e. contusions); AND

History of: Pain in hip or groin at rest or with patient initiated movement (paramedic should not intentionally move joint); AND

Examination: May have externally rotated and/or shortened leg.

## EXCLUSION

1. Patient meets the Trauma Triage Guidelines

2. Patient with hip joint replacement on same side (Pt should be transported to site of original joint replacement surgery. If original site is unknown normal distribution guidelines will apply).

### Education notes:

1. "Isolated" hip fracture: Refers to no other recognized significant injuries.


#### 2. Mechanism:

The intention of the above listed mechanism is to select those patients that are unlikely to have additional injuries (significant trauma mechanism). Although the tool states fall from sitting, lying, standing, this may also include a single step or curb but is meant to exclude more significant falls.

#### 3. Relevant history:

If a patient with a possible hip fracture has an extensive history with one site (i.e. such as post-operative, oncology, dialysis, multiple admissions, or discharged patient), it would be preferable for the patient to be transported to that site (excluding McMaster Children's Hospital or HHS Hamilton General Site).

# Psychiatric Emergency Recognition Tool (HPS)



# Musculoskeletal Injury Recognition Tool (HPS)



For the purposes of Patient Priority System:

Patients with suspected significant orthopedic fractures which might require immediate surgery (excluding hip) by the guidelines below should be transported to the Emergency Departments of St. Joseph's Hospital or Hamilton General Hospital as directed by CACC.

## INCLUSION

Adult patients (≥18) with:

1. Suspected "open" fracture of any limb, OR
2. Severe bony deformity of an injured lower limb

## EXCLUSION

1. Patient's injury is at site of known joint replacement (prosthetic joint), then transport to the Emergency Department to the site where the joint replacement surgery was performed or the Juravinski or St. Joseph's Hospital as directed by CACC.
2. Receiving active oncology treatment at the Juravinski Cancer Clinic, transport to the Juravinski Emergency Department.

## Education notes:

1. If Patient meets the Provincial Trauma Triage Guidelines, then transport to Hamilton General Hospital as directed by CACC.
2. If Patient meets the Possible Hip Fracture Identification Tool, preferentially follow that tool, then transport to the Emergency Department of the Juravinski or St. Joseph's Hospital as directed by CACC.
3. "Open" fracture or compound fracture: Refers to a fracture with an associated wound. This can include circumstances where the bone fragments can be seen protruding through a wound, where there is a large skin defect or even just a small puncture sized wound where the bone may have penetrated the skin but is no longer visible. Any open injury (other than an abrasion) associated with a suspected fracture can be considered a suspected "open" fracture for the purposes of this guideline.
4. The Juravinski Hospital will continue to treat pathological fractures associated with a malignancy
5. All Sites, including the Juravinski Hospital, will continue to manage patients with fractures not requiring immediate surgery, dislocations and soft tissue injuries.

# Ebola Virus Disease (EVD) Screening Recognition Tool



For the purposes of the Patient Priority System:

Patients who are screened as positive (suspected EVD) using the most current Ministry of Health and Long Term Care (MOHLTC) EVD Screening Tool, and who meet specific destination protocol criteria, will be preferentially transported as indicated below:

Adult patient  $\geq 18$  years of age and screened positive for EVD:

- For Decision Priority 1 through 4, follow the current Adult PPS by transporting the patient to the identified destination as per normal practice.
- For Decision Priority 5 through 10, transport the adult patient to the Juravinski Hospital

Pediatric patient  $< 18$  years of age and screened positive for EVD:

- For all Decision Priority criteria follow the current Pediatric PPS by transporting the patient to the identified destination as per normal practice.

Education Notes:

1. When a patient has screened positive for EVD, a patch to notify the receiving facility must be completed by the Paramedics regardless of transport priority.
2. The following hospitals are designated EVD testing sites although the ambulance destination decision will follow the direction above:
  - Juravinski Hospital – Adult patients ( $\geq 18$  years of age)
  - McMaster Children's Hospital – Pediatric patients ( $< 18$  years of age)

# Radio Channel Change Locations



## Hamilton

QEW and Fifty Road=====NIA REG2 COM, contact Hamilton CACC

## London

QEW and Fifty Road=====NIA REG2 COM, contact Hamilton CACC

Hwy 403 and County Road 25 (Middle Townline Road)=====NIA MOH ZN 1, contact London CACC

This is about 15-20 km west of Brantford

## Mississauga

QEW and Fifty Road=====NIA REG2 COM, contact Hamilton CACC

QEW and Hwy 403 (base of Burlington Skyway)=====NIA MOH ZN 1, contact Mississauga CACC

## Toronto

QEW and Fifty Road=====NIA REG2 COM, contact Hamilton CACC

QEW and Hwy 403 (base of Burlington Skyway)=====NIA MOH ZN 1, contact Mississauga CACC

QEW and Hwy 427=====NIA PROV COM, contact Toronto CACC

When returning, the locations for changing back are the same.

If transporting a patient on return to Niagara, switch to NIA TAC 1 at Fifty Road.  
If you are returning empty, switch to NIA North at Fifty Road.

All channels are within the NIA folder and can be found by simply turning the Channel Selector.



Channel Selector

Intro

Airway /  
Breath.

## FAST Sepsis Pre-Alert for GWPS, HPS, and ROWPS

Do you suspect or know there is an infection? If yes, apply ParaHEWS (below)

If ParaHEWS  $\geq 5$ : notify receiving hospital of "Sepsis Pre-Alert" and Apply Capnography

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
ResearchChemical  
ExposureMedical  
Refer.Medic.  
Info.

Contact

Physiological Parameters	3	2	1	0	1	2	3
Heart Rate / Pulse	$\geq 131$	111-130	101-110	51-100	41-50	$< 41$	
Systolic BP	$\geq 201$	171-200		91-170		71-90	$< 71$
Respiratory Rate	$\geq 31$	21-30		14-20		8-13	$< 8$
Temperature (C)		$\geq 39.1$	38.0-39.0	36.1-37.9 (or not available)	35.0-36.0		$< 35$
O <sub>2</sub> Saturation				$\geq 93$	85-92		$< 85$
O <sub>2</sub> Therapy	O <sub>2</sub> via face mask		O <sub>2</sub> via nasal prongs	Room Air			
Change in CNS from Baseline	Not responsive	Pain	Voice	Alert or Usual Self		New Confusion	

[www.sepsis-prealert.ca](http://www.sepsis-prealert.ca)

Destinat.  
Guide.

Destination Guide FAST Sepsis Pre-Alert



# STEMI Protocol Pearls

## Symptoms

### PAIN

Pain can be typical or atypical (but not only non-specific symptoms of dyspnea, nausea, fatigue, etc)

### ACUTE

An acute history of symptoms of < 12 hours



## ECG

### QUALITY

Ensure good quality ECG

- Shave chest
- No moving/talking

### REPEAT

If negative, do serial ECGs

- (1) before treatment
- (2) in ambulance prior to leaving scene
- (3) in ambulance prior to moving into ED

### CAUTION

ECGs can be tricky, rule out mimics  
If not certain, go to closest appropriate ED



## Geography

### 60 MINUTES

Maximum 60 minutes from first medical contact to PCI centre

If you are quicker on scene (eg: 15 minutes), this will allow longer transport time (eg: 45 minutes)



### BOUNDARIES

Know the PCI centres in your area  
CACC may be able to assist

<p>HGH 1-844-832-6830</p> <p>St. Mary's 1-519-653-4074</p>	<p>Brampton 1-416-747-3500,1</p> <p>Southlake 1-905-952-2466</p>
--	--

Trillium  
1-888-493-3568

## Prepare

### CAUTION

Caution with nitro and morphine

Neither of these medications are life-saving in STEMI patients & can cause adverse events

### "PADS ON"

Defibrillation pads are placed on all patients with suspected STEMI



### BE READY

Be familiar with the common complications that can occur:

- dysrhythmias
- pump failure
- cardiac arrest

Be ready to manage them



Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

## Fit2Sit procedure for paramedics arriving at Guelph General Hospital

### Step 1

Enter front Emergency Department doors.

Complete a Fit2Sit assessment/triage form (available in ambulance).

### Step 2

Escort patient to waiting room and provide assistance, if needed, to triage waiting area.

### Step 3

Notify front triage nurse of patient's arrival and give completed Fit2Sit assessment/triage form and patient's Health Card to nurse.

### Step 4

Leave Emergency Department to complete patient's ePCR (electronic patient care reporting) and provide update to dispatch about TOC (transfer of care).





## Fit2Sit criteria



### Inclusion

1. Paramedic determines patient is Fit2Sit
2. Patient age is  $\geq 8$  and  $\leq 70$   
Note: Children must be accompanied by an adult
3. Canadian Triage Acuity Score (CTAS) 3–5
4. Glasgow Coma Scale (GCS) of 15.  
Note: for patients with a baseline below 15, with no change, a family or caregiver must accompany them
5. Vital signs are within defined normal ranges:
  - a. Pulse  $\geq 60$  bpm and  $\leq 120$  bpm
  - b. Blood pressure  $\geq 100$  Systolic and  $\leq 160$
  - c. Respiration  $< 25$ /minute
  - d. Oxygen saturation  $> 92\%$  without supplemental oxygen
6. Patient must be able to ambulate independently (i.e. go to the washroom)

### Exclusion

1. Canadian Triage Acuity Score (CTAS) 1–2
2. Glasgow Coma Scale (GCS) of 14 or lower
3. Patient has clinical condition that requires a stretcher
4. Any symptom relief rendered by paramedics (with exception of ibuprofen or acetaminophen)
5. Patient is a known risk for self-harm or harm to others
6. Patient history of dementia (if no caregiver to stay with patient)
7. Patient complains of chest pain
8. Patient has fever, active vomiting or diarrhea or gross bleeding
9. Overdose or substance abuse

Intro

Airway /  
Breath.

Cardiac /  
Circula.

LOC

Pain/  
Sed./  
Nausea

Proced.

Pall Care /  
Research

Chemical  
Exposure

Medical  
Refer.

Medic.  
Info.

Contact

## List of Mandatory Patch Points

### Medical Cardiac Arrest

- TOR

### Trauma Cardiac Arrest

- Trauma TOR

### Tachydysrhythmia

- amiodarone/lidocaine administration
- adenosine administration if wide complex;
- cardioversion

### IV & Fluid Therapy

- Fluid bolus for hypotensive patients < 12 years of age with suspected DKA

### Analgesia

- morphine/fentanyl administration for patients < 12 years of age
- ketamine < 18 years of age

### Cyanide Exposure

- hydroxocobalamin administration



# Medication Safety Starts with You

When you see the “5Rs” symbol throughout this guidebook, it is a reminder to always confirm:

☐ RIGHT PATIENT

☐ RIGHT DRUG

☐ RIGHT DOSE

☐ RIGHT ROUTE

☐ RIGHT TIME

